The One-Cop-Moves Game on Planar Graphs

Ziyuan Gao * Boting Yang †

Abstract

Cops and Robbers is a vertex-pursuit game played on graphs. In this game, a set of cops and a robber occupy the vertices of the graph and move alternately along the graph's edges with perfect information about each other's positions. If a cop eventually occupies the same vertex as the robber, then the cops win; the robber wins if she can indefinitely evade capture. Aigner and Fromme established that in every connected planar graph, three cops are sufficient to capture a single robber. In this paper, we consider a recently studied variant of the cops and robbers game, alternately called the one-active-cop game, one-cop-moves game or the lazy cops and robbers game, where at most one cop can move during any round. We show that Aigner and Fromme's result does not generalize to this game variant by constructing a connected planar graph on which a robber can indefinitely evade three cops in the one-cop-moves game.

1 Introduction

Cops and Robbers, introduced by Nowakowski and Winkler [16] in 1983 and independently by Quillot [18] in 1978, is a game played on graphs, where a cop tries to capture a robber. The cop is first placed on any vertex of the graph G, after which the robber chooses a starting vertex in G. The cop and robber then move in alternate turns, with the robber moving on odd turns and the cop moving on even turns. A round of the game consists of a robber's turn and the cop's subsequent turn. During every turn, the cop or robber either moves along an edge of G to a neighbouring vertex or stays put on his or her current vertex. Furthermore, both the cop and robber have perfect information about each other's positions at any point in the game. The cop wins the game if he eventually occupies the same vertex as the robber at some moment in the game; the robber wins if she can indefinitely avoid occupying any vertex containing the cop. A winning strategy for the cop on G is a sequence of instructions that, if followed, guarantees that the cop can win any game played on G, regardless of how the robber moves throughout the game. A winning strategy for the robber on G is defined analogously.

Aigner and Fromme [2] studied the original Cops and Robbers game by allowing more than one cop to play; we will henceforth refer to this version of the game as the *cops and robbers* game. They associated to every finite graph G a parameter known as the *(classical) cop number* of G, denoted by c(G), which is the minimum number of cops needed for a cop winning strategy on G, and they showed that the cop number of every connected planar graph is at most 3. Nowakowski

^{*}Department of Mathematics, National University of Singapore, Singapore 119076, Republic of Singapore. Research supported in part by the Singapore Ministry of Education Academic Research Fund Tier 2 grant MOE2016-T2-1-019 / R146-000-234-112. Email: ziyuan84@yahoo.com.

[†]Department of Computer Science, University of Regina, Regina, SK, Canada. Research supported in part by an NSERC Discovery Research Grant, Application No.: RGPIN-2013-261290. Email: Boting.Yang@uregina.ca.

and Winkler [16] gave a characterization of the class of graphs with cop number one. In the same vein, Clarke and MacGillivray [9] characterized the class of graphs with any given cop number. A number of fundamental questions concerning the cop number remain open. In particular, Meyniel's conjecture [6, 7] states that for any graph G of order n, $c(G) = O(\sqrt{n})$. The cops and robbers game has attracted considerable attention from the graph theory community, owing in part to its connections to various graph parameters, as well as the large number of interesting combinatorial problems arising from the study of the cop number. In addition, due to the relative simplicity and naturalness of the cops and robbers game, it has served as a model for studying problems in areas of applied computer science such as artificial intelligence, robotics and the theory of optimal search [8, 11, 15, 21].

This paper examines a variant of the cops and robbers game, known alternately as the oneactive-cop game [17], lazy cops and robbers game [3,4,22] or the one-cop-moves game [25]. The corresponding cop number of a graph G in this game variant is called the *one-cop-moves cop* number of G, and is denoted by $c_1(G)$. One of our motivations for studying the one-cop-moves cop number comes from Meyniel's conjecture: it is hoped that an analogue of Meyniel's conjecture holds in the one-cop-moves game, and it would be easier to prove than the original conjecture (or at least lead to new insights into how Meyniel's conjecture may be proven). The one-cop-moves cop number has been studied for various special families of graphs such as hypercubes [3, 17], generalized hypercubes [20], random graphs [4], Rook's graphs [22], graphs with treewidth at most 2 [24], and Halin graphs [24]. On the other hand, relatively little is known about the behaviour of the one-cop-moves cop number of connected planar graphs. In particular, it is still open at present whether or not there exists an absolute constant k such that $c_1(G) \le k$ for all connected planar graphs G [4, 25]. Instead of attacking this problem directly, one may try to establish lower bounds on $\sup\{c_1(G): G \text{ is a connected planar graph}\}\$ as a stepping stone. Note that the dodecahedron D is a connected planar graph with classical cop number equal to 3 [2]. Since any winning strategy for the robber on D in the cops and robbers game can also be applied to D in the one-cop-moves game, it follows that $c_1(D) \geq 3$, and this immediately gives a lower bound of 3 on $\sup\{c_1(G):G \text{ is a connected planar graph}\}$. To the best of our knowledge, there has hitherto been no improvement on this lower bound. Sullivan, Townsend and Werzanski [22] recently asked whether or not $\sup\{c_1(G): G \text{ is a connected planar graph}\} \geq 4.1$ Many prominent planar graphs have a one-cop-moves cop number of at most 3 (such as the dodecahedron and the truncated icosahedron, known colloquially as the "soccer ball graph") or at most 2 (such as cylindrical grid graphs),² and so the study of such graphs unfortunately does not shed new light on the question. The goal of the present work is to construct a connected planar graph whose structure is specifically designed for a robber to easily evade 3 cops indefinitely, thereby settling the open problem affirmatively. Our graph is a modification of the dodecahedron; for details of the construction and an intuitive explanation of certain features of the graph, see Section 4.

2 Preliminaries

Any unexplained graph terminology is from [23]. The book by Bonato and Nowakowski [6] gives a survey of some proof techniques and important results in the cops and robbers game. All graphs

¹The same question was assigned to Shulang Lei and Rahim Ali in 2012 as their projects when they took the second author's reading course.

²Formal proofs establishing the one-cop-moves cop number of these graphs are usually quite tedious.




Figure 1: Subdivided cube Q'

in this paper are simple, finite and connected. Let G be a graph with n vertices. For any vertex u, a cop λ is said to be k edges away from u iff the distance between the position of λ and u is k; similarly, a vertex v is said to be k edges away from u iff the distance between v and u is k. A path π is defined to be a sequence (v_0, \ldots, v_k) of distinct vertices such that for $0 \le i \le k-1$, v_i and v_{i+1} are adjacent; the length of π is the number of vertices of π minus one. Let u and v be any two distinct vertices of G, and let G be any subgraph of G. If there is a unique shortest path in G from G to G then this path will be denoted by G will be denoted analogously by G will be G will be denoted analogously by G will be G will be denoted analogously by G will be G be a graph with G will be denoted analogously by G will be denoted analogously by G will be denoted by G will be denoted analogously by G will be denoted by G will be denoted analogously by G will be denoted by G will be denoted analogously by G will be denoted by G will be denoted analogously by G will be denoted by G and G will be denoted analogously by G will be denoted by G will be denoted analogously by G will be denoted by G and G and G and G and G are G and G and G are G and G and G are G and G are G and G and G are G and G and G are G are G are G are G and G are G are G are G are G are G and G are G are G are G are G are G are G and G are G are G are G and G are G are G are G are G are G are G and G are G are G

Let $\{\lambda_1,\ldots,\lambda_k\}$ be a set of k cops, and let γ be a robber. The one-cop-moves game is defined as follows. Initially, each of the k cops chooses a starting vertex in G (any two cops may occupy the same vertex); after each cop has chosen his initial position, γ chooses her starting vertex in G. A game configuration (or simply configuration) is a (k+2)-tuple $\langle G, u_1, \ldots, u_k; r \rangle$ such that at the end of some turn of the game, r is the vertex occupied by γ and for $i \in \{1,\ldots,k\}$, u_i is the vertex occupied by λ_i . γ is said to be captured (or caught) if, at any point in the game, γ occupies the same vertex as a cop. The 1-st turn of the game starts after the robber has chosen her starting vertex. During each odd turn $\{1,3,\ldots\}$, the robber γ either stays put or moves to an adjacent vertex, and during each even turn $\{2,4,\ldots\}$, exactly one of the cops moves to an adjacent vertex. For any $i \in \mathbb{N}$, the (2i-1)-st turn and 2i-th turn together constitute the i-th round of the game.

3 The Cops and Robbers Game Versus the One-Cop-Moves Game on Planar Graphs

Before presenting the main result, we show that for planar graphs, the one-cop-moves cop number can in general be larger than the classical cop number. Recall that the cube \mathcal{Q} has domination number 2 and one cop cannot capture a robber on \mathcal{Q} ; so $c(\mathcal{Q}) = c_1(\mathcal{Q}) = 2$. Now let \mathcal{Q}' be the graph obtained by subdividing each edge of \mathcal{Q} with one vertex (see Figure 1). Then we have the following result.

Proposition 3.1. c(Q') = 2 and $c_1(Q') = 3$.

Proof. Let γ denote the robber. We first show that 2 cops can capture γ in the cops and robber game. Initially, we place the cops at vertices 9 and 5. By symmetry, one may assume that γ starts

at one of the following vertices: 1, 2, or 3. The following list shows the possible moves of the game before γ is caught. A triple $\langle p_1, p_2; p_3 \rangle$ denotes the set of positions of the cops and robber at the end of some turn of the game; p_1 and p_2 denote the positions of the first cop and second cop respectively, while p_3 denotes the position of γ . An arrow \rightarrow denotes a transition from one turn to the next turn of the game. The first triple in each sequence denotes the set of positions of the cops and robber at the end of the 1-st turn. It is assumed that whenever the robber is adjacent to a cop at the end of the cops' turn, she will try to escape by moving to an adjacent vertex during the next turn. So c(Q') = 2.

- 1. $\langle 9, 5; 1 \rangle \rightarrow \langle 17, 5; 1 \rangle \rightarrow \langle 17, 5; 8 \rangle \rightarrow \langle 1, 6; 8 \rangle$.
- $2. \ \langle 9,5;1\rangle \rightarrow \langle 17,5;1\rangle \rightarrow \langle 17,5;2\rangle \rightarrow \langle 1,4;2\rangle.$
- 3. $\langle 9, 5; 2 \rangle \rightarrow \langle 17, 4; 2 \rangle \rightarrow \langle 1, 3; 2 \rangle$.
- 4. $\langle 9, 5; 3 \rangle \rightarrow \langle 9, 4; 3 \rangle \rightarrow \langle 9, 4; 2 \rangle \rightarrow \langle 17, 3; 2 \rangle$.
- 5. $\langle 9, 5; 3 \rangle \rightarrow \langle 9, 4; 3 \rangle \rightarrow \langle 9, 4; 18 \rangle \rightarrow \langle 10, 3; 18 \rangle$.

To show that two cops cannot capture the robber on \mathcal{Q}' in the one-cop-moves game, we show that γ can evade capture if she avoids the vertices of degree 2; and if she is forced to move to at least one of these vertices, then she will choose the position that maximizes her total distance from the cops. Assume otherwise. By symmetry, it is enough to show that if γ were eventually caught, then her last position is 8 while the two cops are at vertices (i) 2 and 6 or (ii) 6 and 17.

If (i) holds, then, since 8 cannot be starting position of γ , the previous position of γ must have been either 7 or 1. Again by symmetry, it suffices to assume that the previous position of γ is 7. But if γ is at vertex 7 while the cops are at 2 and 6, then γ would move to vertex 20 on her next turn to maximize her total distance from the cops, a contradiction. For similar reasons, if (ii) holds and γ 's previous position is 7, then she would move to vertex 20 on her next turn.

We next show that $c_1(Q') \leq 3$. Start by placing the cops λ_1, λ_2 and λ_3 at vertices 1, 5 and 11 respectively. Note that γ cannot occupy any one of the following vertices: 2, 17, 8, 10, 18, 12, 4, 6, 19, 1, 5, 11, 3 at the end of the 1-st turn. This leaves the following possible vertices for γ at the end of the 1-st turn: 7, 9, 13, 14, 15, 16, 20. Suppose γ is on vertex 9 after the 1-st turn. γ cannot escape to 1 or to 11 so long as the cops at 1 and 11 stay put. λ_2 then moves to 6. Before λ_2 can block off γ 's last escape path, γ must move to 16. λ_2 then moves to 7. γ must move again before her escape path (9, 16, 15) is cut off by λ_2 , this time moving to 15. λ_3 then moves to 12, preventing γ from escaping along (15, 14, 13). If γ moves back to 16, then λ_1 moves to 17. λ_2 can then move to 20, thus trapping γ in the path (17, 16, 15).

Now suppose γ is on vertex 7 after the 1-st turn. λ_3 then moves to 12. If γ tries to move along (7, 20, 15), then λ_3 moves to block the path (15, 14, 13), first moving to 13. γ also cannot advance along (15, 16, 9) because λ_1 can move to 17 before γ reaches 9. If γ returns along (15, 20, 7), then λ_2 can move to 6, thereby trapping γ along (7, 20, 15).

If γ is on vertex 13, 14, 15, 16 or 20 at the end of the 1-st turn, then the cops can follow up with a winning strategy similar to that in one of the above cases when γ is on vertex 9 or 7 at the end of the 1-st turn.

Having achieved separation between the cops and robbers game and the one-cop-moves game on planar graphs, a question that follows quite naturally is: how large can the gap between c(G) and

 $c_1(G)$ be when G is planar? This question is somewhat more difficult. Although we do not directly address the question in this work, the main result shows that for connected planar graphs, the one-cop-moves cop number can break through the upper bound of 3 for the classical cop number.

Theorem 3.2. There is a connected planar graph \mathcal{D} such that $c_1(\mathcal{D}) \geq 4$.

It may seem excessive to devote an entire paper to a result that only marginally improves the current best lower bound of 3, but the one-cop-moves game appears to be considerably more complex (in terms of possible strategies for the cops and the robber) than the classical game, and as we will explain in Section 4, we obtained the graph \mathcal{D} after attempting a number of simpler variants. We organize the proof of Theorem 3.2 into three main sections. Section 4 details the construction of the planar graph \mathcal{D} with a one-cop-moves cop number of at least 4. Section 5 establishes some preparatory lemmas for the proof that $c_1(\mathcal{D}) \geq 4$. Section 6 describes a winning strategy for a single robber against three cops in the one-cop-moves game played on \mathcal{D} .

4 The Construction of the Planar Graph \mathcal{D}

The basic idea and intuition of the construction. The construction of \mathcal{D} starts with a dodecahedron D.³ This is a fairly natural starting point, given that the dodecahedron has a relatively simple and symmetrical structure, and its classical cop number is already 3. The main idea is to embed a planar graph – the choice of which would favour the robber – into each face of D. A natural strategy for the robber would then be to stay within a "safety zone" in an embedded face of D, and wait until a cop is one edge away from her, upon which the latter would quickly move to the "safety zone" of another face. An earlier idea we considered was to iteratively embed dodecahedrons into each face; however, we were unable to establish that the robber can escape from a face F of a smallest dodecahdron in the graph to another such face when there are 3 cops in F. We also could not provide a straightforward strategy for the robber using a modified version of the icosahedron, which is used in [14]. Another construction we tried was embedding a grid of latitudes and longitudes into the surface of a sphere; this graph, too, did not give an easy strategy for the robber against 3 cops.

The construction of \mathcal{D} . Each vertex of D is called a *corner* of \mathcal{D} . We will add straight line segments on the surface of D to partition each pentagonal face of D into small polygons. For each pentagonal face U of D, we add 48 nested nonintersecting closed pentagonal chains, which are called pentagonal layers, such that each side of a layer is parallel to the corresponding side of U (see Figure 2). Each vertex of a layer is called a corner of that layer. For convenience, the innermost layer is also called the 1-st layer in U and the boundary of U is also called the outermost layer of U or the 49-th layer of U. We add a vertex o in the centre of U and connect it to each corner of U using a straight line segment which passes through the corresponding corners of the 48 inner layers. For each side of the n-th layer $(1 \le n \le 49)$, we add 2n + 1 internal vertices to partition the side path into 2n + 2 edges of equal length (see Figure 3). Add a path of length 2 from the centre vertex o to every vertex of the innermost layer to partition the region inside the 1-st layer into 20 pentagons.

³It is worth noting that a connected planar digraph based on the icosahedron was recently used by Loh and Oh [14] to show that the cop number of directed planar graphs can exceed 3. Similarly, Abrahamsen, Holm, Rotenberg and Wulff-Nilsen [1] recently gave a geometric construction inspired by the dodecahedron to show that a man can escape two lions in a bounded area with rectifiable lakes.




Figure 2: Two innermost and two outermost pentagonal layers of a pentagonal face.

Further, for each pair of consecutive pentagonal layers, say the n-th layer and the (n+1)-st layer $(1 \le n \le 48)$, add paths of length 2 from vertices of the n-th layer to vertices of the (n+1)-st layer such that the region between the two layers is partitioned into 5(2n+2) hexagons and 10 pentagons as illustrated in Figure 2. Let \mathcal{D} be the graph consisting of all vertices and edges currently on the surface of the dodecahedron D (including all added vertices and edges). Since \mathcal{D} is constructed on the surface of a dodecahedron without any edge-crossing, \mathcal{D} must be a planar graph.

Note on terminology. We will treat \mathcal{D} as an embedding of the graph on the surface of D because it is quite convenient and natural to express features of \mathcal{D} in geometric terms. Thus we will often employ geometric terms such as *midpoint*, *parallel*, and *side*; the corresponding graph-theoretic meaning of these terms will be clear from the context. The *distance* between any two vertices u and v in a graph G, denoted $d_G(u, v)$, will always mean the number of edges in a shortest path connecting u and v. Given any $A, B \subseteq V(\mathcal{D})$ and any $v \in V(\mathcal{D})$, define

$$d_{\mathcal{D}}(v,A) = \min\{d_{\mathcal{D}}(v,x) : x \in A\}$$
 and $d_{\mathcal{D}}(A,B) = \min\{d_{\mathcal{D}}(x,y) : x \in A \land y \in B\}.$

By abuse of notation, we will write $d_{\mathcal{D}}(\gamma, v)$ (resp. $d_{\mathcal{D}}(\lambda_i, v)$) to denote the distance between γ and v (resp. between λ_i and v) at the point of consideration. $d_{\mathcal{D}}(\gamma, A)$ and $d_{\mathcal{D}}(\lambda_i, A)$ are defined analogously.

For $n \in \{1, ..., 49\}$, let $L_{U',n}$ denote the *n*-th pentagonal layer of a pentagonal face U', starting from the innermost layer. Define a *side path* of $L_{U',n}$ to be one of the 5 paths of length 2n + 2 connecting two corner vertices of $L_{U',n}$. $L_{U',n}$ will often simply be written as L_n whenever it is clear from the context which pentagonal face L_n belongs to.

The pentagonal faces of \mathcal{D} will be denoted by $U, U_1, U_2, \dots, U_{10}, U_{11}$ (see Figure 4). For $i \in \{1, \dots, 15\}$, B_i will denote a side path of $L_{U',49}$ for some pentagonal face U'. The centre vertex of




Figure 3: A side path L_n ($1 \le n \le 49$) has 2n + 3 vertices and 2n + 2 edges.




Figure 4: 12 pentagonal faces of \mathcal{D} , labelled $U, U_1, \ldots, U_{11}, v_1, \ldots, v_5$ denote the 5 corner vertices of U. The side paths of U are labelled $B_6, B_7, B_8, B_9, B_{10}$; the side paths B_1, B_2, B_3, B_4, B_5 connect U to U_6, U_7, U_8, U_9 and U_{10} respectively. The side paths $B_{11}, B_{12}, B_{13}, B_{14}$ and B_{15} connect U_{11} to U_3, U_4, U_5, U_1 and U_2 respectively. m is the middle vertex of B_1 .

U will be denoted by o, and for $i \in \{1, ..., 11\}$, the centre vertex of U_i will be denoted by o_i . Given a pentagonal face U, we will often abuse notation and write U to denote the subgraph of \mathcal{D} that is embedded on the face U.

For any $n \in \{1, ..., 49\}$, a middle vertex of L_n is a vertex that is n + 1 edges away from two corners of L_n , which are end vertices of some side path of L_n . The middle vertex of a side path B of L_n is the vertex of L_n that lies at the midpoint of B. Given any pentagonal face U, a spoke of U is a path of length 98 connecting a vertex on $L_{U,49}$ and the centre of U.

Let U and U' be any two pentagonal faces of \mathcal{D} . Define $U \cup U'$ to be the subgraph $(V(U) \cup V(U'), E(U) \cup E(U'))$ of \mathcal{D} and $U \cap U'$ to be the subgraph $(V(U) \cap V(U'), E(U) \cap E(U'))$ of \mathcal{D} ; these definitions naturally extend to any finite union or finite intersection of pentagonal faces.

Remark 4.1. The exact number of pentagonal layers in each face of \mathcal{D} is not important so long as it is large enough to allow the robber's winning strategy to be implemented. One could increase the number of pentagonal layers in each face and adjust the robber's strategy accordingly. This will become clearer when we describe the robber's winning strategy in Section 6. One crucial feature of \mathcal{D} is that the distance between the centre of a face U' and the boundary of any pentagonal layer $L_{U',n}$ (for some n with $1 \le n \le 49$) – equal to 2n – is less than the length of a side path of $L_{U',n}$, which is equal to 2n + 2. Intuitively, this particular property of the graph makes it harder for 3 cops to protect the entire boundary of a face while making it comparatively easier for the robber to go from the centre of a face to a vertex on the boundary of the same face.

5 Some Preparatory Lemmas

In this section, we will outline the main types of strategies employed by the robber γ to evade the three cops $\lambda_1, \lambda_2, \lambda_3$. We first state a lemma for determining the distance between any two vertices of a pentagonal face.

Lemma 5.1. Let U be a pentagonal face of \mathcal{D} . Let x be a vertex of L_r and y be a vertex of L_s , where L_r and L_s are pentagonal layers of U and $s \ge r$. Then

$$d_{\mathcal{D}}(x,y) = \min\{2r + 2s, (2s - 2r) + d_{L_r}(w,x)\},\$$

where w is the intersection vertex of L_r and the shortest path between y and the center of U. (See Figure 5 for an illustration.)

Proof. We construct a shortest path from y to x using any given path from y to x. The construction is based on the main ideas of the Floyd-Marshall algorithm [10].

First, consider any path from y to x that passes through o. It may be directly verified that a shortest path from y to o has length 2s while a shortest path from o to x has length 2r. Thus any shortest path from y to x that passes through o has length 2s + 2r.

Second, consider any path from y to x that does not pass through o. Define x' to be the unique vertex on L_s such that $d_{L_s}(x',y) = \min\{d_{L_s}(x'',y) : x'' \text{ lies on } L_s \wedge d_W(x'',x) = s - r\}$.

Suppose $d_{L_s}(x',y) \ge 4s + 4$. Then any shortest path from y to x that does not pass through o covers a distance of at least 4r' + 4 along a layer $L_{r'}$ for some least r'. Since any path along $L_{r'}$ of length at least 4r' + 4 can be replaced by a shorter path of length 4r' passing through o, the length of any path from y to x is at least 2r + 2s.




Figure 5: Distance between x and y in \mathcal{D}

Now suppose that $d_{L_s}(x',y) \leq 4s+3$. Observe that any path π that starts at a vertex z in a pentagonal layer L_{r_1} , goes to a neighbouring layer L' – which includes L_{r_1+1} if $r_1 \leq 48$, L_{r_1-1} if $r_1 \geq 2$, and the 48th layer of a neighbouring face if $r_1 = 49$, and then passes along L', covering a distance equal to at most twice the length of a side path of L' when traversing L', before returning to a vertex z' in L_{r_1} , may be replaced with a path π' that goes directly from z to z' along L_{r_1} such that the length of π' is not more than that of π . Thus any shortest path from y to x that does not pass through o may be replaced with one that goes from L_s to L_r , passing in succession the intermediate pentagonal layers L_i with r < i < s (and possibly passing along each layer). Next, observe that for any $r_2 \geq 2$, any path θ that starts at a vertex z in L_{r_2} , passes along L_{r_2} , and then goes directly to a vertex z' in L_{r_2-1} , may be replaced with a path θ' that starts at z, goes directly to L_{r_2-1} in 2 rounds, and then passes along L_{r_2-1} before ending at z'; in addition, the length of θ' does not exceed that of θ . Applying this observation iteratively and combining it with the earlier observation that any shortest path from y to x that does not pass through o may be replaced with one that starts by going directly from L_s to L_r in s-r steps, one obtains a path from y to x that starts from y, goes directly to a vertex w belonging to L_r in 2s-2r rounds, and then slides along the shortest path in L_r from w to x before ending at x; furthermore, the length of this path is not more than that of any other path from y to x that does not pass through o.

The following observation will often be used implicitly to simplify subsequent arguments.

Lemma 5.2. Suppose that γ is currently at vertex a_1 of \mathcal{D} and a cop λ is currently at vertex u. Suppose γ starts moving towards vertex a_{n+1} via the path $(a_1, a_2, \ldots, a_{n+1})$. Then, by the 2n-th turn of the game (starting at the turn when γ moves from a_1 to a_2), γ can reach a_{n+1} without being caught by λ if $d_{\mathcal{D}}(u, a_{n+1}) > n$.

Proof. Suppose that λ catches γ on the 2k-th turn of the game for some $k \in \{1, \ldots, n\}$. It follows that $d_{\mathcal{D}}(u, a_{k+1}) \leq k$. Since $d_{\mathcal{D}}(a_{k+1}, a_{n+1}) \leq n - k$, one has $d_{\mathcal{D}}(u, a_{n+1}) \leq d_{\mathcal{D}}(u, a_{k+1}) + d_{\mathcal{D}}(a_{k+1}, a_{n+1}) \leq k + (n - k) = n$. This is a contradiction.

Suppose that the robber γ currently occupies o. Consider any set $A \subseteq V(\mathcal{D})$ of vertices. For every $v \in A$, if there is a cop λ such that the current distance between λ and v is less than $d_{\mathcal{D}}(o, v)$, then by Lemma 5.2, λ can capture γ if γ tries moving to v (assuming that γ starts the game).

Corollary 5.3. Suppose that γ is currently at the centre o of a pentagonal face U and there is a centre $o' \neq o$ such that $d_{\mathcal{D}}(o, o') < d_{\mathcal{D}}(\lambda_j, o')$ for all $j \in \{1, 2, 3\}$. Then γ can reach o' without being caught.

The following lemma is a direct consequence of Lemma 5.1.

Lemma 5.4. Suppose a cop λ lies at a vertex u in a pentagonal face U of \mathcal{D} and is not at the centre of U. Let A be the set of 5 corners of $L_{U,49}$. If, for some set $A' \subseteq A$, $d_U(u,v) \leq 98$ whenever $v \in A'$, then $|A'| \leq 2$. Furthermore, if there are two corners v', v'' of $L_{U,49}$ such that $d_U(u,v') \leq 98$ and $d_U(u,v'') \leq 98$, then $d_U(v',v'') = 100$. Let M be the set of 5 middle vertices of $L_{U,49}$. If, for some set $M' \subseteq M$, $d_U(u,v) \leq 98$ whenever $v \in M'$, then $|M'| \leq 2$.

The next technical lemma will be used to devise an evasion tactic for γ in a set of game configurations. More generally, the sort of tactic described in the proof of this lemma will often be used by γ to escape to the centre of a pentagonal face. It may be described informally as follows. γ starts from the centre of a pentagonal face U and she first tries to move to the centre of a neighbouring face, say U'. Then at least one cop (say λ_1) will be forced to protect the centre of U'. Just before λ_1 can catch γ in U', γ deviates from her original path towards the centre of U' and moves towards the centre of yet another neighbouring face, say U'', such that γ is closer to the centre of U'' than λ_1 is. Since at most one cop can move during any round, the speed of the remaining two cops (λ_2 and λ_3) will be reduced as λ_1 is chasing γ . Thus all three cops will be sufficiently far away from the centre of U'' during the round when γ deviates from her original path, and this will allow γ to successfully reach the centre of U''.

Lemma 5.5. Suppose the one-cop-moves game played on \mathcal{D} starts on γ 's turn with the following configuration (illustrated in Figure 6). γ lies at the centre o of the pentagonal face U and the 3 cops lie in U. Let u_1, u_2 and u_3 denote the vertices currently occupied by λ_1, λ_2 and λ_3 respectively. Let m' be any middle vertex of $L_{U,49}$, and let B be the side path of $L_{U,49}$ containing m'. Let p' be any vertex in B that is 1 edge away from m'. Suppose that $d_{\mathcal{D}}(u_2, m') \geq 99$ and $d_{\mathcal{D}}(u_3, m') \geq 99$ (resp. $d_{\mathcal{D}}(u_2, p') \geq 99$ and $d_{\mathcal{D}}(u_3, p') \geq 99$). Suppose that $d_{\mathcal{D}}(u_1, o) = 1$ and $d_{\mathcal{D}}(u_i, B) + d_{\mathcal{D}}(u_j, B) \geq 104$ (resp. $d_{\mathcal{D}}(u_i, B) + d_{\mathcal{D}}(u_j, B) \geq 110$) for all distinct $i, j \in \{1, 2, 3\}$. Assume that $d_{\mathcal{D}}(u_1, m') \geq 98$ (resp. $d_{\mathcal{D}}(u_1, p') \geq 98$) and both $d_{\mathcal{D}}(u_2, o) \geq 2$ and $d_{\mathcal{D}}(u_3, o) \geq 2$ hold. Then γ can reach the centre of a pentagonal face at some point after the first round of the game without being caught.

Proof. Suppose that $d_{\mathcal{D}}(u_2, m') \geq 99$ and $d_{\mathcal{D}}(u_3, m') \geq 99$, $d_{\mathcal{D}}(u_i, B) + d_{\mathcal{D}}(u_j, B) \geq 104$ for all distinct $i, j \in \{1, 2, 3\}$ and $d_{\mathcal{D}}(u_1, m') \geq 98$ (the proof for the other case is entirely similar). The proof of this lemma will be explained with the aid of Figure 4.

Suppose that $m'=m_2$, so that $B=B_7$. γ begins by moving towards m_2 , traversing the middle vertices of the side paths of $L_1, L_2, \ldots, L_{U,49}$ parallel to B_7 . Note that if λ_1 moves 1 step into L_1 during the first round of the game, then γ can simply move back to o during her next turn without being caught. Now suppose that λ_1 does not move during the first round of the game. Then γ can safely reach m_2 in 98 rounds. After the 98-th round of the game, the total distance travelled by λ_1, λ_2 and λ_3 is at most 98. Suppose that γ reaches m_2 in the 98-th round. Consider the following case distinction.

Case (a): For each $j \in \{1, 2, 3\}$, the distance between λ_j and U_2 at the end of the 98-th round is at least 1. By Lemma 5.2, γ can reach o_2 in another 98 rounds without being caught.




Figure 6: The relative positions of the cops and γ .

Case (b): At least one of λ_1, λ_2 and λ_3 occupies a vertex of U_2 at the end of the 98-th round of the game. Note that since $d_{\mathcal{D}}(u_i, B_7) + d_{\mathcal{D}}(u_j, B_7) \geq 104$ for all distinct $i, j \in \{1, 2, 3\}$, it follows that at least 104 rounds are needed for a minimum of two cops to reach B_7 , and therefore at most one of λ_1, λ_2 and λ_3 can occupy a vertex of U_2 at the end of the 98-th round of the game.

Let λ_{α} be the first cop that reaches U_2 and s be the first vertex of U_2 that λ_{α} reaches as γ is moving from o to m_2 . Without loss of generality, assume that s lies on B_7 . Note that s cannot be m_2 (since γ can safely reach m_2 in 98 rounds), and therefore either $d_{\mathcal{D}}(s, B_1) > d_{\mathcal{D}}(s, B_2)$ or $d_{\mathcal{D}}(s, B_1) < d_{\mathcal{D}}(s, B_2)$ holds. Assume that $d_{\mathcal{D}}(s, B_1) > d_{\mathcal{D}}(s, B_2)$. For each $j \in \{1, 2, 3\}$, let $\ell_j = d_{\mathcal{D}}(u_j, B_7)$. Note that $\ell_1 \geq 97$ and for each fixed $j \in \{2, 3\}$, if u'_j is a vertex on B_7 such that $d_{\mathcal{D}}(u_j, u'_j) = \ell_j$, then $\ell_j = d_{\mathcal{D}}(u_j, u'_j) \geq d_{\mathcal{D}}(u_j, m_2) - d_{\mathcal{D}}(m_2, u'_j) \geq 99 - 50 = 49$. Let k be the total distance travelled by λ_{α} between the 1-st and the 98-th round.

Case (b.1): $k \ge \ell_{\alpha} + 46$. Since $46 + \ell_{\alpha} \le k \le 98$, it holds that $\ell_{\alpha} \le 52$ and therefore $\alpha \in \{2, 3\}$. Without loss of generality, assume that $\alpha = 2$.

 γ moves along the path $m_2 \xrightarrow{B_7} v_1 \xrightarrow{B_1} m$ (where m is the midpoint of B_1). Since s lies on B_7 and $d_{\mathcal{D}}(s, B_1) > d_{\mathcal{D}}(s, B_2)$ by assumption, an application of Lemma 5.1 shows that the shortest path from s to m passes through m_2 . As γ can reach m_2 in 98 rounds but λ_2 needs at least 99 rounds to reach m_2 , it follows that λ_2 cannot catch γ before or during the round when γ reaches m. Furthermore, for $j \in \{1,3\}$, the distance between λ_j and m at the end of the 98-th round is at least $\ell_j + 50 - 98 + k \ge (\ell_j + \ell_2) - 48 + 46 \ge 102$ (since λ_j could have moved at most 98 - k steps between the 1-st and the 98-th round and $d_{\mathcal{D}}(B_7, m) = 50$). Since the distance between γ and m at the end of the 98-th round is 100, it follows from Lemma 5.2 that for $j \in \{1,3\}$, λ_j cannot catch γ either before or during the round when γ reaches m.

If λ_2 moves at most 96 steps between the 99-th round and the 198-th round, then γ can reach o_6 via $m \xrightarrow{B_1} q_1 \xrightarrow{S} o_6$, where S is the spoke connecting q_1 and o_6 . If λ_2 moves at least 97 steps between the 99-th round and the 198-th round, then any $\lambda \in \{\lambda_1, \lambda_3\}$ can move at most 3 steps between

⁴The phrase "between the *m*-th round of the game and the *n*-th round of the game" will always mean "between the *m*-th round of the game and the *n*th-round of the game *inclusive*" (unless explicitly stated otherwise).

the 99-th round and the 198-th round. Note that if λ_2 is in $U_1 \cup U_6$ but not in U_2 at the end of the 198-th round, then γ can safely move from m to o_2 in another 98 rounds. It will therefore be assumed that at the end of the 198-th round, λ_2 is at least 196 edges away from w. γ now starts moving from m to o_1 (via the spoke connecting m and o_1).

We claim that for some appropriate choice of r, γ can either reach o_1 or move to a vertex of L_r and thence to w without being caught via the roundabout path $m \rightsquigarrow s_1 \leadsto s_2 \leadsto t \leadsto w$ shown in Figure 7.

First, note that between the 1-st and the 98-th round, the total distance travelled by λ_1 and λ_3 is at most $98 - k \le 98 - \ell_2 - 46 \le 3$. Thus the total distance travelled by λ_1 and λ_3 between the 1-st and the 198-th round is at most 6, so that when γ is at m, the distance between w and the cop that is nearest to w (say λ_3) is at least 190. A direct calculation gives that the length of the path $m \rightsquigarrow s_1 \leadsto s_2 \leadsto t \leadsto w$ is 2(98 - 2r) + (r+1) + (2r+2) = 199 - r, and so by Lemma 5.2, choosing any $r \ge 10$ ensures that λ_3 will not be able to catch γ before or during the round when γ reaches w. In particular, for any $r \ge 10$, λ_3 will not be able to catch γ during the round when γ reaches s_1 . Now suppose that $r \ge 10$. If, between the 198-th round and the round when γ reaches s_1 , s_2 at least 7 turns, then s_3 will be closer to s_3 than any other cop just after the round when s_3 reaches s_4 , and therefore s_3 can reach s_4 without being caught.

Suppose, on the other hand, that λ_3 skips no more than 6 turns as γ is moving from m to s_1 . Then, just after the round when γ reaches s_1 , λ_2 must be at least 141 edges away from w. Thus by choosing r so that the distance from s_1 to w (via the path highlighted in Figure 7) is less than 141 steps, γ can reach w without being caught by λ_2 . Therefore one requires 3r + 3 + 98 - 2r = 101 + r < 141, or r < 40. Fixing any r in the range of 10 to 39 (inclusive) establishes the claim. After reaching w, γ can safely reach o_{10} in another 98 rounds by moving along the spoke connecting w and o_{10} .

Case (b.2): $k \le \ell_{\alpha} + 45$.

 γ adopts a winning strategy similar to that in Case (b.1), this time moving towards o_2 . As in Case (b.1), we claim that for some appropriate choice of r, γ can either reach o_2 without being caught or move to q_1 and thence to o_6 without being caught via the path $m_2 \rightsquigarrow s'_1 \leadsto s'_2 \leadsto t' \leadsto q_1$ highlighted in Figure 8. We will again assume that $\alpha = 2$; it will become clear below that the following winning strategy for γ also works for $\alpha \in \{1,3\}$. r is defined according to Algorithm 1.

We briefly explain how Algorithm 1 works. γ moves successively through $w_1, w_2, \ldots, w_{k-\ell_2+1}$ until at least one of the following occurs: (i) she reaches some w_i such that the total number of turns j_i that λ_2 skips between the round when γ is at w_0 (:= m_2) and the round when γ is at w_i is exactly equal to i-1, or (ii) she reaches $w_{k-\ell_2+1}$. At this stage, Algorithm 1 breaks out of the loop. Let w_ℓ be the last vertex that γ reaches just before Algorithm 1 stops; then r and s'_1 are defined to be r_ℓ and w_ℓ respectively.

Note that $4 \le r_1 \le 49$. Set $j_0 = 0$. A straightforward induction shows that for all $i \in \{1, \ldots, \ell\}$, $j_{i-1} \ge i-1$. We show by induction on $i \in \{0, 1, \ldots, \ell\}$ that γ can safely get from w_0 (:= m_2) to w_i in $98 - 2r_i$ rounds. To show that γ can safely reach w_i , it suffices to show that λ_2 cannot catch γ before or during the round when γ reaches w_i .

The case i=0 was established earlier. For the inductive step, suppose that for some $i \in \{1, ..., \ell\}$, γ can safely reach w_{i-1} in $98 - 2r_{i-1}$ rounds. We first calculate a lower bound for the distance

Algorithm 1: Algorithm for computing r

```
1 r_0 \longleftarrow 49;
 \mathbf{2} \ w_0 \longleftarrow m_2;
 3 for i = 1 to k - \ell_2 + 1 do
        r_i \longleftarrow k - \ell_2 + 5 - i;
 5
        w_i \leftarrow middle vertex of the side path of L_{r_i} parallel to B_7;
 6
        r \longleftarrow r_i;
        move \gamma from w_{i-1} to w_i in 2(r_{i-1} - r_i) rounds;
 7
        j_i \leftarrow number of turns that \lambda_2 skips between the round when \gamma is at
 8
         w_0 and the round when \gamma is at w_i;
        if j_i = i - 1 then
 9
         break;
10
        end
11
12 end
13 return r;
```

between λ_2 and w_i at the end of the round when γ reaches w_{i-1} . Now, any path from s to w_i that passes through o_2 has length at least 99. On the other hand, the path from s to w_i that starts by going directly to L_{r_i} in $98 - 2r_i$ rounds and then passing along the side path of L_{r_i} parallel to B_7 until w_i is reached has length at most $(98 - 2r_i) + (r_i + 1) = 99 - r_i \le 99 - 4 = 95$. Therefore no shortest path from s to w_i passes through o_2 . Hence by Lemma 5.1, a shortest path from s to w_i starts by going directly to L_{r_i} in $98 - 2r_i$ rounds, and then passing along the side path of L_{r_i} parallel to m_2 until w_i is reached. Denote this shortest path from s to w_i by P. Observe that the distance between s and m_2 is at least $99 - \ell'_2$, where $\ell'_2 = d_{\mathcal{D}}(u_2, s)$. Thus the shortest distance between w_i and the first vertex of P on L_{r_i} is either $99 - \ell'_2$ or $r_i + 1$. Note that λ_2 moves a distance of at most $k - \ell'_2$ between the round he reaches s and the 98-th round. In addition, λ_2 moves at most $98 - 2r_{i-1} - j_{i-1}$ steps between the round when γ is at w_0 and the round when γ is at w_i . It follows that at the end of the round when γ reaches w_{i-1} , the distance between λ_2 and w_i is at least

$$\min\{((98 - 2r_i) + (99 - \ell_i')) - (k - \ell_2') - (98 - 2r_{i-1} - j_{i-1}), ((98 - 2r_i) + (r_i + 1)) - (k - \ell_2') - (98 - 2r_{i-1} - j_{i-1})\}$$

$$= \min\{2(r_{i-1} - r_i) + (99 - k + j_{i-1}), 2(r_{i-1} - r_i) + (r_i + 1 - k + \ell_2' + j_{i-1})\}.$$

Note that γ needs $2(r_{i-1} - r_i)$ rounds to get from w_{i-1} to w_i . Since $k \leq 98$, $99 - k + j_{i-1} \geq 1$. Similarly,

$$r_i + 1 - k + \ell'_2 + j_{i-1} = k - \ell_2 + 5 - i + 1 - k + \ell'_2 + \underbrace{j_{i-1}}_{\geq i-1}$$

 $\geq \underbrace{(\ell'_2 - \ell_2)}_{\geq 0} + 5 \geq 5$.

Consequently, γ can move from w_{i-1} to w_i in $2(r_{i-1} - r_i)$ rounds without being caught by λ_2 , and this completes the inductive step.

If $j_{\ell} > k - \ell_2$, then after reaching s'_1 , γ continues moving towards o_2 until she reaches o_2 in another 2r rounds. Suppose $j_{\ell} \leq k - \ell_2$. It can be directly verified that in this case, the condition to break




Figure 7: The escape path of γ in Case (b.1).




Figure 8: The escape path of γ in Case (b.2).

out of the loop in Algorithm 1 will eventually be satisfied, and that $r=k-\ell_2+4-j_\ell$. γ now moves along the path $p' \leadsto s'_1 \leadsto s'_2 \leadsto q_1$ highlighted in Figure 8. Note that between the 1-st round and the round when γ reaches s'_1 , λ_1 and λ_3 could have moved a total of at most $j_\ell+(98-k)$ steps. Suppose that λ_3 chases γ for the duration of γ 's movement from s'_1 to q_1 . During the round when γ is at s'_1 , the distance between λ_3 and q_1 is at least $(100+\ell_3)-(98-k+j_\ell)=\ell_2+\ell_3+2+(k-\ell_2-j_\ell)$. The length of the path $s'_1 \leadsto s'_2 \leadsto t' \leadsto q_1$ is $101+r=101+(k-\ell_2+4-j_\ell)=105+k-\ell_2-j_\ell$. Since $(\ell_2+\ell_3+2+(k-\ell_2-j_\ell))-(105+(k-\ell_2-j_\ell))=\ell_2+\ell_3-103\geq 1$, γ can reach q_1 without being caught by λ_3 . One can show in an analogous way that γ can reach q_1 without being caught by λ_1 . If λ_2 chases γ by moving along L_r (or by any other path that does not pass through o_2), then, since γ can safely get from m_2 to s'_1 in 98-2r rounds, λ_2 cannot catch γ before or during the round when γ reaches q_1 . Suppose λ_2 chases γ by first moving to o_2 and then to q_1 . The number of rounds required by λ_2 to move from his position when γ is at s'_1 to q_1 by taking a path passing through o_2 is at least $196-((98-2r-j_\ell)+(k-\ell_2))=98+2r-k+\ell_2+j_\ell=98+2(k-\ell_2+4-j_\ell)-k+\ell_2+j_\ell=106+k-\ell_2-j_\ell$. Thus γ can reach q_1 without being caught by λ_2 . After reaching q_1 , γ can safely reach o_6 in another 98 rounds.

The following lemma will establish a winning strategy for γ in another specific game configuration. As in Lemma 5.5, γ 's strategy in Lemma 5.6 exploits the condition that at most one cop can move

during any round. Roughly speaking, the strategy works as follows: when γ is at a corner v, she attempts to lure a cop into a face U' containing v by moving to a neighbour of v in U'. If no cop is in U' at the end of the next turn, then γ can safely reach the centre of U'; otherwise, γ safely moves back to v during the next round and repeats the same strategy used during the preceding round. Lemma 5.6 shows that it is advantageous for γ to occupy a corner, and this fact underlies γ 's strategy as described in Section 6.

Lemma 5.6. Suppose γ is currently at a vertex $v \in V(U \cap U')$, and it is γ 's turn. Suppose λ_1 is at some vertex w of $U \cup U'$ such that $d_{\mathcal{D}}(v, w) \geq 1$, $d_{\mathcal{D}}(\lambda_2, U \cup U') \geq 2$ and $d_{\mathcal{D}}(\lambda_3, U \cup U') \geq 2$. Then γ can either (i) reach the centre of U or U' without being caught, or (ii) oscillate infinitely often between v and one of its neighbours.

Proof. We prove this lemma by induction on the odd turns of the game. Assume that the 1-st turn starts on γ 's turn. Inductively, suppose that at the start of the (2n-1)-st turn of the game (for some $n \geq 1$), γ is at a vertex v of $U \cap U'$, λ_1 is at some vertex w_n of $U \cup U'$ such that $d_{\mathcal{D}}(v, w_n) \geq 1$, the distance between the position of every cop (other than λ_1) and $U \cup U'$ is at least 2, and it is currently γ 's turn. Without loss of generality, assume that w_n belongs to U. γ then moves to a vertex v' in U' such that v' is adjacent to v and the distance between v' and the centre of U' is 97. If λ_1 does not move towards the centre of U' during the 2n-th turn or if w_n belongs to $V(U) \setminus V(U')$, then, since $d_{\mathcal{D}}(v, w_n) \geq 1$ and the distance between every cop (other than λ_1) and $U \cup U'$ is at least 2, γ can continue moving safely towards the centre of U', reaching this vertex in another 97 rounds. On the other hand, if λ_1 does move towards the centre of U' on the 2n-th turn and w_n is in $U \cap U'$, then γ moves back to v during the (2n+1)-st turn without being caught. Note that in this case, at the start of the (2n+2)-nd turn, λ_1 is at a vertex w' of $U \cup U'$ such that $d_{\mathcal{D}}(v,w') \geq 2$, and the distance between every other cop and $U \cup U'$ is still at least 2. If λ_1 is not in $U \cap U'$ at the end of the (2n+2)-nd turn, then γ can safely reach the centre of U using another 98 turns. If λ_1 is in $U \cap U'$ at the end of the (2n+2)-nd turn, then at the start of the (2n+3)-rd turn, λ_1 is at some vertex w_{n+1} of $U \cap U'$ such that $d_{\mathcal{D}}(v, w_{n+1}) \geq 1$ while each of the other two cops is 2 edges away from $U \cap U'$. This completes the induction step.

The next lemma is the analogue of Lemma 5.6 when γ lies at the intersection of 3 pentagonal faces.

Lemma 5.7. Suppose γ is currently at a corner $v \in V(U \cap U' \cap U'')$, and it is γ 's turn. Suppose moreover that there are at most 2 cops, say λ_1 and λ_2 , lying in $U \cup U' \cup U''$, and $d_{\mathcal{D}}(\lambda_1, \gamma) \geq 1$, $d_{\mathcal{D}}(\lambda_2, \gamma) \geq 2$ and $d_{\mathcal{D}}(\lambda_3, U \cup U' \cup U'') \geq 2$. Then γ can either (i) reach the centre of U, or the centre of U', or the centre of U'' without being caught, or (ii) oscillate infinitely often between v and one of its neighbours.

Proof. For any two faces U and U' of \mathcal{D} , define $U \setminus U'$ to be the subgraph of \mathcal{D} induced by $V(U) \setminus V(U') \cup (V(U') \setminus V(U)) \cup (V(U') \setminus V(U))$. Like the proof of Lemma 5.6, we use induction on the odd turns of the game. Inductively, suppose that at the start of the (2n-1)-st turn of the game (for some $n \geq 1$), γ is at vertex v and the three cops are positioned according to one of the following arrangements: (1) one cop (say λ_2) is in $(U \triangle U') \triangle U''$ and is at a distance of at least 2 from γ , one cop (say λ_1) is in $((U \cap U') \setminus U'') \cup ((U' \cap U'') \setminus U) \cup ((U \cap U'') \setminus U')$ and is at a distance of at least 1 from γ , and the remaining cop (say λ_3) is at a distance of at least 2 from $U \cup U' \cup U''$; (2) both λ_2 and λ_1 are in $((U \cap U') \setminus U'') \cup ((U' \cap U'') \setminus U) \cup ((U \cap U'') \setminus U')$; the distance between λ_2 and γ is at least 2, and the

distance between λ_3 and $U \cup U' \cup U''$ is at least 2; (3) both λ_2 and λ_1 are in $(U \triangle U') \triangle U''$ and are each at a distance of at least 1 from γ , and the distance between λ_3 and $U \cup U' \cup U''$ is at least 1.

Suppose (1) holds. Without loss of generality, assume that λ_2 is in $U \setminus (U' \cup U'')$ and is at a distance of at least 2 from γ , and λ_1 is in $(U' \cap U'') \setminus U$ and is at a distance of at least 1 from γ . γ then takes 1 step towards the centre of U'. Suppose λ_1 does not move towards the centre of U' on the 2n-th turn. Then, since both λ_2 and λ_3 are at a distance of at least 99 from the centre of U' at the start of the (2n-1)-st turn, γ can safely reach the centre of U'. Now suppose λ_1 moves towards the centre of U' on the 2n-th turn of the game. γ then moves back to v during the (2n+1)-st turn. If λ_1 does not return to a vertex of $(U' \cap U'') \setminus U$ during the (2n+2)-nd turn of the game, then γ can safely reach the centre of U'' in another 98 rounds. If λ_1 returns to a vertex of $(U' \cap U'') \setminus U$ during the (2n+2)-nd turn, then scenario (1) is repeated at the start of the (2n+3)-rd turn.

Suppose (2) holds. Without loss of generality, assume that λ_2 is in $(U \cap U') \setminus U''$ and λ_1 is in $(U' \cap U'') \setminus U$. γ then moves towards the centre of U. If, during the 2n-th turn, λ_2 does not move towards the centre of U, then γ can safely reach the centre of U in another 98 rounds. If λ_2 moves towards the centre of U during the 2n-th turn, then γ returns to v during the (2n+1)-st turn. If λ_2 does not move back to a vertex of $(U \cap U') \setminus U''$ during the (2n+2)-nd turn, then either (1) or (3) holds at the start of the (2n+3)-rd turn. If λ_2 does move back to a vertex of $(U \cap U') \setminus U''$ during the (2n+2)-nd turn, then scenario (2) is repeated at the start of the (2n+3)-rd turn.

Suppose (3) holds. Without loss of generality, suppose λ_2 is in $U \setminus (U' \cup U'')$ and λ_1 is in $U' \setminus (U \cup U'')$. γ can then reach safely the centre of U'' in 98 rounds. This completes the induction step.

6 The Robber's Winning Strategy: Proof of Theorem 3.2

We begin with a high-level description of γ 's winning strategy; see Algorithm 2.

Algorithm 2: High-level strategy for γ

- 1 γ picks the centre of a pentagonal face that is free of cops. Let U be this face.
- $_{2}$ γ stays at the centre o of U until there is exactly one cop that is 1 edge away from γ .
- 3 γ does one of the following depending on the cops' positions and strategy (details will be given in Cases (A), (B) or (C) below; see Sections 6.2, 6.3 and 6.4): (i) she moves to the centre of a pentagonal face U', which may or may not be U, without being caught at the end of a round, or (ii) she oscillates back and forth along an edge for the rest of the game without being caught.
- 4 If, in Step 3, γ does (i), then set $U \leftarrow U'$ and go back to Step 2.

Since there are 12 pentagonal faces but only 3 cops, Step 1 of Algorithm 2 can be readily achieved. Let U denote the pentagonal face whose centre o is currently occupied by γ . The precise winning strategy for γ in Step 3 will depend on the relative positions of the cops when exactly one cop is 1 edge away from γ .⁵ The details of this phase of γ 's winning strategy will be described in three cases: (A) when three cops lie in U; (B) when exactly one cop lies in U; (C) when exactly two cops lie in U. These cases reflect three possible strategies for the cops: all three cops may try to

⁵In order to reduce the number of cases in our proof, we choose to let the robber wait until a cop is exactly one edge away from her; by symmetrical considerations, it would suffice to assume that when the robber starts moving away from her current position o, there is exactly one cop occupying one of only three possible vertices adjacent to o (refer to p_1, p_2, p_3 in Figure 2).

Algorithm 3: A strategy for γ when γ is at a corner

- 1 Suppose γ is at a corner v and it is γ 's turn. Let U, U' and U'' be the faces containing v.
- 2 If there is a centre o' such that $d_{\mathcal{D}}(v, o') \leq d_{\mathcal{D}}(\lambda_i, o')$ for all $i \in \{1, 2, 3\}$, then move γ from v to o' in $d_{\mathcal{D}}(v, o')$ rounds and stop the procedure.
- **3** If there are two distinct faces $U_i, U_j \in \{U, U', U''\}$ and there is one cop (say λ_1) such that $d_{\mathcal{D}}(\lambda_1, v) \geq 1$, $d_{\mathcal{D}}(\lambda_2, U_i \cup U_j) \geq 2$ and $d_{\mathcal{D}}(\lambda_3, U_i \cup U_j) \geq 2$, then apply Lemma 5.6.
- 4 If there are at most two cops (say λ_1 and λ_2) in $U \cup U' \cup U''$ such that $d_{\mathcal{D}}(\lambda_1, v) \geq 1$ and $d_{\mathcal{D}}(\lambda_2, v) \geq 2$, while the third cop λ_3 satisfies $d_{\mathcal{D}}(\lambda_3, U \cup U' \cup U'') \geq 2$, then apply Lemma 5.7.
- **5** Else, move γ to another corner and go back to Step 1.⁶

encircle γ , or one cop may try to chase γ while the remaining two cops guard the neighbouring faces of U, or two cops may try to encircle γ while the remaining cop guards the neighbouring faces of U.

We will frequently use the following general subroutine in γ 's strategy (details depend on the individual cases considered).

6.1 Assumptions and notation for Cases (A), (B) and (C)

It will be assumed that the starting game configurations in Cases (A), (B) and (C) below occur during the first round of the game (so that in what follows, for any $n \ge 1$, the "n-th round of the game" refers to the n-th round of the game after the given initial game configuration) and that γ starts each round. That is, the inputs of Algorithms 4, 5 and 6 will be the initial game configurations.

Now suppose that in each starting game configuration of Cases (A), (B) and (C), γ lies at the centre o of U, λ_1 is exactly 1 edge away from γ , and it is γ 's turn to move. By symmetrical considerations, it suffices to assume that λ_1 is positioned at p_1 , p_2 or p_3 as shown in Figure 2. If λ_1 moves away from o during the second turn of the game (so that λ_1 is 2 edges away from o at the end of the first round), then γ can simply return to o during the second round. Thus in our analysis of γ 's strategies in Cases (A), (B) and (C), it will be assumed that λ_1 either stays still or moves to o during the first round of the game. Let u_1, u_2 and u_3 be the starting vertices occupied by λ_1, λ_2 and λ_3 respectively.

6.2 Case (A): U contains three cops when γ lies at o and $d_{\mathcal{D}}(\lambda_1, o) = 1$

Note that there is at most one corner v' of $L_{U,49}$ such that $d_{\mathcal{D}}(u_1, v') \leq 98$. Let v_1, v_2, v_3, v_4, v_5 be the 5 corner vertices of $L_{U,49}$, labelled clockwise, and m_1, m_2, m_3, m_4, m_5 be the 5 middle vertices of $L_{U,49}$, also labelled clockwise. The vertex p is 1 edge away from m_4 and lies between m_4 and v_3 , and the vertex q is 1 edge away from m_5 and lies between m_5 and v_4 (see Figure 4). We summarise γ 's strategy in Algorithm 4; the detailed analysis of this algorithm can be found in Appendix A. As was mentioned earlier, every corner of \mathcal{D} is a strategic location for γ , and so γ will generally try to reach a corner if no cop is protecting it. To give an example of how Algorithm 4 works, suppose the starting configuration $\langle \mathcal{D}, p_1, m_1, m_3; o \rangle$ (see Figures 2 and 4) is fed to Algorithm 4. By Step 3 of Algorithm 4, Lemma 5.5 will be applied. According to the strategy given in the proof of Lemma 5.5, γ will first move to m_4 in 98 rounds. If no cop is in U_4 at the end of the 98-th round, then γ

⁶If the conditions in Steps 2, 3 and 4 are not satisfied, then we use a strategy similar to the one given in the proof of Lemma 5.5 to move γ from a corner to another corner.

Algorithm 4: The Robber's Strategy for Case (A)

- 1 Suppose the current game configuration is $\langle \mathcal{D}, u_1, u_2, u_3; o \rangle$, where o is the centre of face U, $\{u_1, u_2, u_3\} \subset V(U), u_1 \in \{p_1, p_2, p_3\}$ (see Figure 2), $d_{\mathcal{D}}(u_2, o) \geq 2$ and $d_{\mathcal{D}}(u_3, o) \geq 2$.
- 2 If there is a corner v of $L_{U,49}$ such that $d_{\mathcal{D}}(u_1,v) \geq 98$, $d_{\mathcal{D}}(u_2,v) \geq 99$ and $d_{\mathcal{D}}(u_3,v) \geq 99$, since λ_1 cannot move towards v in the 1-st round, we can move γ from o to v in 98 rounds and then call Algorithm 3.
- 3 If there does not exist a corner v of $L_{U,49}$ satisfying the condition in Step 2, then apply Lemma 5.5.

can safely reach o_4 in another 98 rounds; otherwise, a straightforward calculation shows that at the end of the 98-th round, λ_2 cannot be in U_4 while at most one of $\{\lambda_1, \lambda_3\}$ is in U_4 . If either λ_1 or λ_3 is in U_4 at the end of the 98-th round, then γ continues moving towards o_4 until she reaches $L_{U_4,r}$ for some r depending on the relative movements of the cops (refer to Algorithm 1); at this point, she either moves safely to o_4 or deviates from her original path towards o_4 and moves to either q_4 and then to o_9 or to q_3 and then to o_8 .

6.3 Case (B): U contains only λ_1 when γ lies at o and $d_{\mathcal{D}}(\lambda_1, o) = 1$

We split γ 's strategy into two main subcases: either (i) there is a corner of $L_{U,49}$ that γ can reach in 98 rounds without being caught, or (ii) for every corner v of $L_{U,49}$, at least one of the following holds: (a) at least one of $\{\lambda_2, \lambda_3\}$ is at a distance of at most 98 from v, or (b) λ_1 is at a distance of 97 from v. Each subcase is further broken into cases depending on the relative initial positions of the cops. The specific strategies used by γ in each subcase are similar to those in Case (A) but the details are more tedious. γ 's strategy in the present case is summarised in Algorithm 5 (see Appendix B for the detailed analysis).

Algorithm 5: The Robber's Strategy for Case (B)

- 1 Suppose the current game configuration is $\langle \mathcal{D}, u_1, u_2, u_3; o \rangle$, where o is the centre of face U, $\{u_2, u_3\} \cap V(U) = \emptyset$ and $u_1 \in \{p_1, p_2, p_3\}$ (see Figures 2 and 4).
- 2 If there is a corner $v \in \{v_1, v_2, v_3, v_4, v_5\}$ of $L_{U,49}$ such that $d_{\mathcal{D}}(u_1, v) \geq 98$, $d_{\mathcal{D}}(u_2, v) \geq 99$, and $d_{\mathcal{D}}(u_3, v) \geq 99$, w.l.o.g., assume that $v = v_1$, then set $F \longleftarrow U_{10} \cup U_6 \cup U_1 \cup U_2 \cup U_7$ and we have the following cases.
 - (2.1) If u_2 and u_3 are in F, then, depending on the position of u_1 , move γ from o to one of $\{v_1, v_3, v_4\}$, then call Algorithm 3.
 - (2.2) If neither u_2 nor u_3 is in F, then move γ from o to v_1 in 98 rounds and then call Algorithm 3.
 - (2.3) If only one of u_2 and u_3 is in F, e.g., u_2 is in U_1 and u_3 is in U_3 (other cases are trivial or similar), then we have two subcases.
 - (2.3.1) If $d_{\mathcal{D}}(u_3, F) \geq 99$, then move γ from o to one of $\{v_1, v_5\}$ in 98 rounds and then call Algorithm
 - 3. (2.3.2) If $d_{\mathcal{D}}(u_3, F) < 99$, then either move γ from o to v_1 in 98 rounds and then call Algorithm 3 or move γ from o to a vertex between v_4 and m_4 or m_5 in 98 rounds and then move to one of $\{o_4, o_5\}$, or one of $\{q_4, z_2, t_9\}$ and then call Algorithm 3.
- 3 Suppose that there does not exist a corner $v \in \{v_1, v_2, v_3, v_4, v_5\}$ of $L_{U,49}$ such that $d_{\mathcal{D}}(u_1, v) \geq 98$, $d_{\mathcal{D}}(u_2, v) \geq 99$, and $d_{\mathcal{D}}(u_3, v) \geq 99$. W.l.o.g., assume that u_2 is in U_1 while u_3 is in U_3 (other cases are trivial or similar). From the condition, we have $d_{\mathcal{D}}(u_1, v_4) = 97$. So γ can reach one of $\{m_2, m_4, m_5\}$ and then apply a strategy similar to the proof of Lemma 5.5.

6.4 Case (C): U contains exactly two cops when γ lies at o and $d_{\mathcal{D}}(\lambda_1, o) = 1$

Without loss of generality, assume that λ_3 is in U and λ_2 is not in U. As in Case (A), we divide γ 's winning strategy into two subcases depending on whether or not γ can safely reach a corner of $L_{U,49}$ in 98 rounds. γ 's winning strategy is outlined in Algorithm 6. The detailed analysis is given in Appendix C.

Algorithm 6: The Robber's Strategy for Case (C)

- Suppose the current game configuration is $\langle \mathcal{D}, u_1, u_2, u_3; o \rangle$, where o is the centre of face $U, u_3 \in V(U)$, $u_2 \notin V(U)$ and $u_1 \in \{p_1, p_2, p_3\}$ (see Figures 2 and 4).
- 2 If there is a corner v of $L_{U,49}$ such that $d_{\mathcal{D}}(u_1,v) \geq 98$, $d_{\mathcal{D}}(u_2,v) \geq 99$ and $d_{\mathcal{D}}(u_3,v) \geq 99$, w.l.o.g., assume that $v = v_1$, then set $F \leftarrow U_{10} \cup U_6 \cup U_1 \cup U_2 \cup U_7$ and we have the following cases. (2.1) If u_2 is in F, then we have two subcases.
 - (2.1.1) If $d_{\mathcal{D}}(u_2, v_5) \leq 98$, then apply one of the following strategies: (i) move γ from o to one of $\{v_1, v_2\}$ in 98 rounds and then call Algorithm 3, (ii) move γ from o to m_5 in 98 rounds and then apply a strategy similar to the proof of Lemma 5.5, or (iii) move γ from o to p in 98 rounds and then apply Lemma 5.5.
 - (2.1.2) If $d_{\mathcal{D}}(u_2, v_5) \geq 99$, then move γ from o to one of $\{v_1, v_2, v_3, v_5\}$ in 98 rounds and then call Algorithm 3.
 - (2.2) If u_2 is not in F, then either move γ from o to v_1 in 98 rounds and then call Algorithm 3 or move γ from o to p in 98 rounds and then apply a strategy similar to the proof of Lemma 5.5.
- 3 Suppose that there does not exist a corner v of $L_{U,49}$ such that $d_{\mathcal{D}}(u_1,v) \geq 98$, $d_{\mathcal{D}}(u_2,v) \geq 99$ and $d_{\mathcal{D}}(u_3,v) \geq 99$. W.l.o.g., assume that u_2 is in U_1 . From the condition we have $d_{\mathcal{D}}(u_1,v_4) = 97$. So γ can reach one of $\{m_2, m_4, m_5\}$ and then apply a strategy similar to the proof of Lemma 5.5.

From the strategy described in the above algorithms, we know that at least 4 cops are necessary for capturing γ on \mathcal{D} .

7 Concluding Remarks

The present work established separation between the cops and robbers game and the one-cop-moves game on planar graphs by exhibiting a connected planar graph whose one-cop-moves cop number exceeds the largest possible classical cop number of connected planar graphs. We believe that this result represents an important first step towards understanding the behaviour of the one-cop-moves cop number of planar graphs. It is hoped, moreover, that some of the proof techniques used in this work could be applied more generally to the one-cop-moves game played on any planar graph.

This work did not prove any upper bound for the one-cop-moves cop number of \mathcal{D} ; nonetheless, we conjecture that 4 cops are sufficient for catching the robber on \mathcal{D} . A characterization of k-copwin graphs for the one-cop-moves game is given in [24]. A more general characterization for cops and robbers games is also given in [5]. If we want to use these characterizations to show if \mathcal{D} is 4-copwin in the one-cop-moves game, basically we need to check almost every pair (v, S), where v is a vertex of \mathcal{D} and S is a multisubset of 4 vertices of \mathcal{D} . However, the graph \mathcal{D} has 302,762 vertices, and so it has 302762⁴ multisubsets of 4 vertices. Thus the methods in [5,24] need about 1.2×10^{23} steps to check if the relation is complete, which is highly impractical for \mathcal{D} .

It should also be noted that the Planar Separator Theorem of Lipton and Tarjan [13] may be applied to show that the one-cop-moves cop number of every connected planar graph with n vertices is at most $O(\sqrt{n})$ (the proof is essentially the same as that in the case of planar directed graphs; see [14, Theorem 4.1]). It may be asked whether or not the robber has a simpler winning strategy on

 \mathcal{D} than that presented in this paper. We have tried a number of different approaches to the problem, but all of them led to new difficulties. For example, one might suggest reducing Case (B) to Case (C) by allowing a single cop to chase the robber in a pentagonal face U until a second cop arrives in U. However, such a strategy would generate new cases to consider since the relative positions of the robber and cop in U just before a second cop reaches U may vary quite widely. One reason it is not quite so easy to design a winning strategy for the robber on \mathcal{D} is that a key lemma of Aigner and Fromme in the cops and robbers game [2] – that a single cop can protect all the vertices of any shortest path P, in the sense that after a bounded number of rounds, if the robber ever moves onto a vertex of P, she will be captured by the cop – carries over to the one-cop-moves game.

The question of whether or not there exists a constant k such that $c_1(G) \leq k$ for all connected planar graphs G [25] remains open. It is tempting to conjecture that such an absolute constant does exist.

References

- [1] M. Abrahamsen, J. Holm, E. Rotenberg and C. Wulff-Nilsen. Best laid plans of lions and men. Preprint. https://arxiv.org/pdf/1703.03687.pdf.
- [2] M. Aigner and M. Fromme. A game of cops and robbers. Discrete Applied Mathematics 8(1984):1–12.
- [3] D. Bal, A. Bonato, W. B. Kinnersley and P. Pralat. Lazy cops and robbers on hypercubes. *Combinatorics Probability and Computing* 24(6):829–837, 2015.
- [4] D. Bal, A. Bonato, W. B. Kinnersley and P. Pralat. Lazy cops and robbers played on random graphs and graphs on surfaces. *International Journal of Combinatorics* 7(4):627–642, 2016.
- [5] A. Bonato, G. MacGillivray, Characterizations and algorithms for generalized cops and robbers games. *Contributions to Discrete Mathematics* 12(1):110–122, 2017.
- [6] A. Bonato and R. Nowakowski. The Game of Cops and Robbers on Graphs. American Mathematical Society, Providence, 2011.
- [7] A. Bonato (2016). Conjectures on cops and robbers. In *Graph Theory: Favorite Conjectures* and *Open Problems 1* (pages 31–42). Springer International Publishing.
- [8] T. H. Chung, G. A. Hollinger and V. Isler. Search and pursuit-evasion in mobile robotics. *Autonomous Robots* 31(4):299–316, 2011.
- [9] N. E. Clarke and G. MacGillivray. Characterizations of k-copwin graphs. *Discrete Mathematics* 312(2012):1421–1425.
- [10] T. H. Cormen, C. E. Leiserson and R. L. Rivest. Introduction to Algorithms. Cambridge, Massachusetts: The MIT Press, 2009.
- [11] A. Isaza, J. Lu, V. Bulitko and R. Greiner. A cover-based approach to multi-agent moving target pursuit. *Proceedings of the 4th Conference on Artificial Intelligence and Interactive Digital Entertainment*, pages 54–59, 2008.

- [12] W. Kinnersley. Cops and robbers is EXPTIME-complete. *Journal of Combinatorial Theory Series B* 111(2015):201–220.
- [13] R. Lipton and R. Tarjan. A separator theorem for planar graphs. SIAM Journal on Applied Mathematics, 36(1979):177–189.
- [14] P. Loh and S. Oh. Cops and robbers on planar directed graphs. *Journal of Graph Theory*, to appear.
- [15] C. Moldenhauer and N. R. Sturtevant. Evaluating strategies for running from the cops. Proceedings of the 21st International Joint Conference on Artificial intelligence, IJCAI'09, pages 584–589, 2009.
- [16] R. Nowakowski and P. Winkler. Vertex to vertex pursuit in a graph. *Discrete Mathematics* 43(1983):235–239.
- [17] D. Offner and K. Okajian. Variations of Cops and Robber on the hypercube. *Australasian Journal of Combinatorics* 59(2):229–250, 2014.
- [18] A. Quilliot. Jeux et pointes fixes sur les graphes. Thèse de 3ème cycle, Université de Paris VI, 1978, pages 131–145.
- [19] A. Quilliot. A short note about pursuit games played on a graph with a given genus. *Journal of Combinatorial Theory Series B* 38(1985): 89–92.
- [20] K. A. Sim, T. S. Tan and K. B. Wong. Lazy cops and robbers on generalized hypercubes. *Discrete Mathematics* 340(2017):1693–1704.
- [21] F. Simard, M. Morin, C. Quimper, F. Laviolette and J. Desharnais. Bounding an optimal search path with a game of cop and robber on graphs. *Principles and Practice of Constraint Programming: 21st International Conference, CP 2015, Cork, Ireland, August 31 September 4, 2015*, Proceedings. Springer International Publishing:403–418.
- [22] B. W. Sullivan, N. Townsend and M. Werzanski. The 3 × 3 rooks graph is the unique smallest graph with lazy cop number 3. Preprint. https://arxiv.org/abs/1606.08485.
- [23] D. B. West. Introduction to Graph Theory. Prentice Hall, 2000.
- [24] B. Yang. The One-Cop-Moves Game on Graphs of Small Treewidth. Submitted.
- [25] B. Yang and W. Hamilton. The optimal capture time of the one-cop-moves game. *Theoretical Computer Science* 588(2015):96–113.

A Detailed analysis of Algorithm 4 for Case (A)

Case (1) There is at least one corner v' of $L_{U,49}$ such that $d_{\mathcal{D}}(v',u') \geq 99$ for all $u' \in \{u_2,u_3\}$ and $d_{\mathcal{D}}(v',u_1) \geq 98$. First, assume that $d_{\mathcal{D}}(v_1,u') \geq 99$ for all $u' \in \{u_2,u_3\}$; since $u_1 \in \{p_1,p_2,p_3\}$, it holds that $d_{\mathcal{D}}(v_1,u_1) \geq 99$. γ first moves to v_1 in 98 rounds. If, at the end of the 98-th round, both




Figure 9: Initial positions of λ_1 and γ

 λ_2 and λ_3 are at least 101 edges away from q_1 , then γ can safely reach q_1 in another 100 rounds and Lemma 5.7 may then be applied to $U_1 \cup U_2 \cup U_6$.

Suppose, on the other hand, that either λ_2 or λ_3 is at most 100 edges away from q_1 at the end of the 98-th round. Without loss of generality, assume that at the end of the 98-th round, λ_2 is in U_1 and is at most 100 edges away from q_1 . This implies that λ_3 could have moved at most 1 step between the 1-st round and the 98-th round, and so λ_3 is at most 1 edge closer to o_2 than γ is at the end of the 98-th round. γ now starts moving towards o_2 . If λ_3 skips more than 1 turn as γ is approaching o_2 , then γ can safely reach o_2 ; else, γ continues moving towards o_2 until she reaches vertex t'' on $L_{U_2,23}$, as shown in Figure 10. γ then moves along the path $t'' \rightsquigarrow t'''$ highlighted in Figure 10. Note that the length of the path $t'' \rightsquigarrow t'''$ is $2 \cdot 23 + 2 = 48$, while λ_2 is at least $98 - 2 \cdot 23 = 52$ edges away from t''' and λ_3 is at least $4 \cdot 23 - 1 = 91$ edges away from t''' when γ is at t''. It follows that γ can safely reach t'''. Furthermore, suppose that between the round when γ is at t'' and the round when γ is at t''', λ_2 moves i steps and λ_3 moves j steps. Since $i+j \leq 2 \cdot 23 + 2 = 48$, at least one of the following holds: (i) $i \le 24$; (ii) $j \le 24$. If (i) holds, then, since λ_2 is at least $97 - i \ge 73$ edges away from q_1 (and λ_3 is even further away from q_1) while γ is $98 - 2 \cdot 23 = 52$ edges away from q_1 when γ is at t''', γ can safely move to q_1 ; Lemma 5.7 may then be applied to $U_1 \cup U_2 \cup U_6$. If (ii) holds, then γ moves along the path $t''' \rightsquigarrow t'''' \rightsquigarrow t_{12}$ highlighted in Figure 10. Note that the length of the path $t''' \rightsquigarrow t'''' \rightsquigarrow t_{12}$ is 100, whereas λ_3 is at least $97 + 2 \cdot 23 - j \ge 119$ edges away from t_{12} (and λ_2 is even further away from t_{12}) when γ is at t'''. Consequently, γ can safely move to t_{12} in another 100 rounds after reaching t'''. Upon reaching t_{12} , either γ may safely move to o_7 in another 98 rounds, or (if λ_3 uses up at least $99 - j \ge 75$ turns to move towards U_7 as γ starts moving from t''' to t'''') Lemma 5.7 may be applied to $U_2 \cup U_6 \cup U_7$.

Case (1'): For some $v'' \in \{v_2, v_3, v_4, v_5\}$, $d_{\mathcal{D}}(v'', u') \geq 99$ for all $u' \in \{u_2, u_3\}$ and $d_{\mathcal{D}}(v'', u_1) \geq 99$. Notice that γ 's strategy in Case (1) still applies (with an appropriate transformation of vertices; for example, if $v'' = v_2$, then we apply the mapping $v_2 \to v_1, v_1 \to v_5, v_5 \to v_4, v_4 \to v_3, v_3 \to v_2$, and extend this mapping so as to obtain an automorphism of \mathcal{D}).

Case (2): For every corner v' of $L_{U,49}$, there is some $u' \in \{u_2, u_3\}$ such that $d_{\mathcal{D}}(u', v') \leq 98$ or $d_{\mathcal{D}}(u_1, v') \leq 97$ (or both inequalities hold). Without loss of generality, assume that $d_{\mathcal{D}}(u_1, v_4) = 97$, $d_{\mathcal{D}}(u_2, v_1) \leq 98$, $d_{\mathcal{D}}(u_2, v_5) \leq 98$, $d_{\mathcal{D}}(u_3, v_2) \leq 98$ and $d_{\mathcal{D}}(u_3, v_3) \leq 98$. Recall that q is the vertex of $L_{U,49}$ that is one edge away from m_5 and p is the vertex of $L_{U,49}$ that is one edge away from m_4 (as shown in Figure 4). Note that by Lemma 5.1, the condition imposed on the positions of λ_2 and λ_3 ,




Figure 10: The escape path of γ in Case (A.1).

and the fact that neither λ_2 nor λ_3 is at o, it holds that $d_{\mathcal{D}}(u_2, B_9) \geq 99$ and $d_{\mathcal{D}}(u_3, B_{10}) \geq 99$.

Case (2.1): $d_{\mathcal{D}}(p, u_3) \geq 99$. As was observed earlier, $d_{\mathcal{D}}(u_2, B_9) \geq 99$. We show that $d_{\mathcal{D}}(u_3, B_9) \geq 50$. Take any $x \in V(B_9)$. If x lies between v_4 and m_4 inclusive, then $d_{\mathcal{D}}(u_3, x) \geq d_{\mathcal{D}}(x, v_2) - d_{\mathcal{D}}(u_3, v_2) \geq 150 - 98 = 52$. If x lies between p and v_3 inclusive, then $d_{\mathcal{D}}(u_3, x) \geq d_{\mathcal{D}}(p, u_3) - d_{\mathcal{D}}(p, x) \geq 99 - 49 = 50$. Since $d_{\mathcal{D}}(u_2, B_9) + d_{\mathcal{D}}(u_3, B_9) \geq 149$, Lemma 5.5 shows that γ can reach the centre of a pentagonal face.

Case (2.2): $d_{\mathcal{D}}(q, u_2) \geq 99$. One can establish in a way similar to that used in Case (2.1) the inequality $d_{\mathcal{D}}(u_2, B_{10}) \geq 48$, so that $d_{\mathcal{D}}(u_2, B_{10}) + d_{\mathcal{D}}(u_3, B_{10}) \geq 147$. An application of Lemma 5.5 then gives the required result.

Case (2.3): $d_{\mathcal{D}}(p, u_3) \leq 98$ and $d_{\mathcal{D}}(q, u_2) \leq 98$. Then, by Lemma 5.1 and the fact that $\{u_2, u_3\} \cap \{o\} = \emptyset$, $d_{\mathcal{D}}(u_2, m_2) \geq 99$ and $d_{\mathcal{D}}(u_3, m_2) \geq 99$. For any $x \in V(B_7)$, $d_{\mathcal{D}}(u_2, x) \geq d_{\mathcal{D}}(x, q) - d_{\mathcal{D}}(q, u_2) \geq 151 - 98 = 53$ and $d_{\mathcal{D}}(u_3, x) \geq d_{\mathcal{D}}(x, p) - d_{\mathcal{D}}(u_3, p) \geq 149 - 98 = 51$. Thus $d_{\mathcal{D}}(u_2, B_7) + d_{\mathcal{D}}(u_3, B_7) \geq 104$, and so one may conclude from Lemma 5.5 that γ can move to m_2 and safely reach the centre of a pentagonal face.

B Detailed analysis of Algorithm 5 for Case (B)

Case (1): There is at least one corner $v_i \in \{v_1, v_2, v_3, v_4, v_5\}$ of $L_{U,49}$ such that $d_{\mathcal{D}}(v_i, u') \geq 99$ for all $u' \in \{u_1, u_2, u_3\}$. We consider the case i = 1; the proofs for the cases i = 2, 3, 4, 5 are similar. Define $F := U_{10} \cup U_6 \cup U_1 \cup U_2 \cup U_7$.

Case (1.1): Both u_2 and u_3 are in F.

Case (1.1.1): $d_{\mathcal{D}}(u_2, U_1 \cup U_2) + d_{\mathcal{D}}(u_3, U_1 \cup U_2) \ge 100$. First, suppose that at least one of u_2 and u_3 belongs to $V(U_1) \cup V(U_2)$. Without loss of generality, assume that $u_2 \in V(U_1) \cup V(U_2)$. Then $d_{\mathcal{D}}(u_3, U_1 \cup U_2) \ge 100$.

Suppose $u_2 \in V(B_1)$. γ first moves to v_3 in 98 rounds. If λ_1 does not reach U_4 by the end of the 98-th round, then γ can safely reach o_4 in another 98 rounds. If λ_1 does reach U_4 by the end of the




Figure 11: The escape path of γ in Case (B.1.1.2.1).

98-th round, using up at least 97 turns in the process, then after γ reaches v_3 , Lemma 5.7 may be applied to $U \cup U_3 \cup U_4$.

Suppose $u_2 \notin V(B_1)$. γ first moves to v_1 in 98 rounds. If λ_2 does not move to B_1 as γ is moving to v_1 , then B_1 does not contain any cop at the end of the 98-th round and at least one of U_1 and U_2 , say U_i , does not contain any cop at the end of the 98-th round; thus γ can safely reach o_i in another 98 rounds. If λ_2 does move to B_1 as γ is moving to v_1 , using up at least 1 turn in the process, then both λ_3 and λ_1 are at least 2 edges away from $U_1 \cup U_2$ at the end of the 98-th round, and so Lemma 5.6 may be applied to $U_1 \cup U_2$.

Second, suppose that neither u_2 nor u_3 belongs to $V(U_1) \cup V(U_2)$. γ then moves to v_1 in 98 rounds. Since $d_{\mathcal{D}}(u_i, U_1 \cup U_2) + d_{\mathcal{D}}(u_j, U_1 \cup U_2) \geq 100$ for any distinct $i, j \in \{1, 2, 3\}$, at least two of the cops are more than 1 edge away from $U_1 \cup U_2$ at the end of the 98-th round. Hence after γ reaches v_1 , Lemma 5.6 may be applied to $U_1 \cup U_2$.

Case (1.1.2): $d_{\mathcal{D}}(u_2, U_1 \cup U_2) + d_{\mathcal{D}}(u_3, U_1 \cup U_2) \leq 99.$

Case (1.1.2.1): At least one of u_2 and u_3 is more than 100 edges away from $U_3 \cup U_4 \cup U_8$. Without loss of generality, assume that u_2 is at least 100 edges away from $U_3 \cup U_4 \cup U_8$. γ first moves to v_3 in 98 rounds. If λ_1 does not reach U_4 by the end of the 98-th round, then γ can safely reach o_4 in another 98 rounds. Suppose λ_1 does reach U_4 by the end of the 98-th round, using up at least 97 turns in the process. Then, since u_3 is at most 99 edges away from $U_1 \cup U_2$, λ_3 must be at least 100 edges away from q_3 at the end of the 98-th round. γ now continues moving towards o_4 until she reaches $L_{U_4,48}$ as shown in Figure 11; she then uses 98 turns to move from p'' to p'''. Suppose that as γ is moving from p'' to p''', λ_3 moves at most 97 steps. Then γ can safely move from p''' to q_3 in another 2 rounds; after γ reaches q_3 , Lemma 5.7 may be applied to $U_3 \cup U_4 \cup U_8$. Suppose that as γ is moving from p'' to p''', λ_3 moves exactly 98 steps. Then γ moves from p''' to p'''' and then to t_9 along the path highlighted in Figure 11. After reaching t_9 , γ may then safely move to o_9 in another 98 rounds.

Case (1.1.2.2): Both u_2 and u_3 are at most 100 edges away from $U_3 \cup U_4 \cup U_8$. If $d_{\mathcal{D}}(u_2, B_1) \geq 99$, then γ moves to v_1 and then to o_1 in 196 rounds.

Suppose $d_{\mathcal{D}}(u_2, B_1) \leq 98$. Since $d_{\mathcal{D}}(u_2, U_3 \cup U_4 \cup U_8) \leq 100$, $d_{\mathcal{D}}(u_2, B_1) = 98$. If $d_{\mathcal{D}}(u_3, U_1 \cup U_2) \geq 2$, γ moves to v_1 ; after reaching v_1 , either γ can safely move to one of o_1 and o_2 , or Lemma 5.6 may be applied to $U_1 \cup U_2$. If $d_{\mathcal{D}}(u_3, U_1 \cup U_2) \leq 1$, γ moves to v_3 ; then, arguing as in Case (1.1.2.1), either




Figure 12: The escape path of γ in Case (B.1.2).

 γ may safely reach o_4 in another 98 rounds, or γ may safely reach q_3 and then apply the strategy in Lemma 5.7.

Case (1.2): Neither u_2 nor u_3 is in F. Note that $d_{\mathcal{D}}(u_2, q_1) \geq 197$ and $d_{\mathcal{D}}(u_3, q_1) \geq 197$. γ first moves to v_1 in 98 rounds. If at least one of u_2 and u_3 is not in $U_3 \cup U_5$, Lemma 5.6 may be applied to $U_1 \cup U_2$ after γ reaches v_1 . We will therefore assume that u_2 is in U_5 and u_3 is in U_3 (the remaining cases can be dealt with in a very similar way).

If both λ_2 and λ_3 are at least 101 edges away from q_1 at the end of the 98-th round, then γ can safely reach q_1 in another 100 rounds. After γ reaches q_1 , either γ may safely move to o_6 in another 98 rounds or Lemma 5.7 may be applied to $U_1 \cup U_2 \cup U_6$.

Suppose, on the other hand, that at least one of λ_2 and λ_3 is at most 100 edges away from q_1 at the end of the 98-th round. Without loss of generality, assume that λ_2 is at most 100 edges away from q_1 at the end of the 98-th round. If λ_2 is at most 99 edges away from q_1 at the end of the 98-th round, then neither λ_1 nor λ_3 could have moved between the 1-st and the 98-th round, and therefore λ_2 can safely reach o_2 in another 98 rounds. Suppose that λ_2 is exactly 100 edges away from q_1 and λ_3 is in U_2 at the end of the 98-th round. γ then starts moving towards o_2 until she reaches vertex x' on $L_{U_2,23}$ as shown in Figure 12. She then moves from x' to x'' in 48 rounds (see Figure 12). If, after γ reaches x'', λ_2 is still at least 53 edges away from q_1 (meaning that λ_2 did not move during 1 round as γ went from x' to x''), γ can safely reach q_1 in another 52 rounds and Lemma 5.7 may then be applied to $U_1 \cup U_2 \cup U_6$. Suppose that after γ reaches x'', λ_2 is 52 edges away from q_1 . γ then continues moving along the path highlighted in Figure 12 until she reaches x'''.

Again, if λ_3 skips at least one turn as γ is moving from x'' to x''', then γ can safely move to x'''' and then move to t_{12} ; she may then apply the strategy in Lemma 5.7 to $U_2 \cup U_6 \cup U_7$.

On the other hand, if λ_3 does not skip any turn as γ is moving from x'' to x''', then γ continues moving along the path highlighted in Figure 12 until she reaches m_7 . If, just after γ reaches m_7 , λ_2 is still at least 1 edge away from U_6 , then γ can safely reach o_6 . If λ_2 is in U_6 just after γ reaches m_7 , then λ_3 must still be at least 52 edges away from t_{12} when γ is at m_7 . γ may thus safely move from m_7 to t_{12} in 50 rounds, and then apply the strategy in Lemma 5.7 to $U_2 \cup U_6 \cup U_7$.




Figure 13: An escape path of γ in Case (B.1.3.2.1.1.1).

Case (1.3): Exactly one of u_2 and u_3 is in F. We will assume that u_2 is in U_1 and u_3 is in U_3 (the other cases are trivial or similar).

Case (1.3.1): $d_{\mathcal{D}}(u_3, F) \geq 99$. First, suppose $u_2 \in V(B_1)$. Then γ can safely move to v_5 in 98 rounds. If, at the end of the 98-th round, λ_1 is not in U_5 , then γ can safely reach o_5 in another 98 rounds. Suppose λ_1 does reach U_5 by the end of the 98-th round, using up at least 97 turns in the process. γ can then safely reach q_5 by moving along B_5 . After γ reaches q_5 , Lemma 5.6 may be applied to $U_1 \cup U_5 \cup U_{10}$.

Second, suppose $u_2 \notin V(B_1)$. γ first moves to v_1 in 98 rounds. Suppose λ_2 does not move as γ is moving to v_1 . Then, since $d_{\mathcal{D}}(u_2, U_1 \cup U_2) \geq 99$ and $d_{\mathcal{D}}(u_3, F) \geq 99$, no cop occupies a vertex belonging to $V(B_1)$ at the end of the 98-th round; furthermore, both λ_2 and λ_3 are at least 1 edge away from $U_1 \cup U_2$ at the end of the 98-th round. Hence, after reaching v_1 , γ can safely reach either o_1 or o_2 .

Now suppose λ_2 uses up at least 1 turn as γ is moving to v_1 . Then both λ_2 and λ_3 must be at least 2 edges away from $U_1 \cup U_2$ at the end of the 98-th round. γ may now apply the strategy in Lemma 5.6 to $U_1 \cup U_2$.

Case (1.3.2): $d_{\mathcal{D}}(u_3, F) \leq 98$.

Case (1.3.2.1): $d_{\mathcal{D}}(u_3, v_3) \geq 12$. γ begins moving towards m_4 . We further distinguish two cases.

Case (1.3.2.1.1): λ_1 moves at least 47 steps as γ is moving towards m_4 . Suppose that as γ is approaching m_4 , λ_1 moves z steps for some $z \geq 47$. γ then continues moving until she reaches m_4 in 98 rounds. Note that λ_2 and λ_3 can move a total of at most 51 steps between the turn γ moves away from o and the turn after γ reaches m_4 . So λ_3 is at most 39 edges closer to o_4 than γ is after γ reaches m_4 . We may assume that at least one of λ_1, λ_3 reaches U_4 just after γ reaches m_4 (otherwise, γ can safely reach o_4 in another 98 rounds).

Case (1.3.2.1.1.1): λ_1 reaches U_4 before λ_3 . Note that λ_3 is still at least 11 edges away from U_4 just after γ reaches m_4 . γ starts moving towards o_4 until she reaches $L_{U_4,4}$; γ then moves along the path highlighted in Figure 13. An argument very similar to those used in earlier cases shows that either γ can move to o_8 without being caught after reaching r''', or γ can continue moving until she safely reaches t_9 , at which point Lemma 5.7 may be applied to $U_4 \cup U_8 \cup U_9$.




Figure 14: An escape path of γ in Case (B.1.3.2.1.1.2).

Case (1.3.2.1.1.2): λ_3 reaches U_4 before λ_1 . Suppose that λ_3 is ℓ edges closer to o_4 than γ is during the turn after γ reaches m_4 . Note that $\ell \leq 39$. γ starts by moving towards o_4 . Suppose that as γ is approaching o_4 , λ_3 skips j turns. If $j > \ell$ then γ can safely reach o_4 . So assume that $j \leq \ell$. γ continues moving towards o_4 until she reaches $L_{U_4,4+\ell-j}$. She then moves along the path highlighted in Figure 14. One can verify that after reaching q_3 , either γ can safely reach o_8 in another 98 rounds, or Lemma 5.7 may be applied to $U_3 \cup U_4 \cup U_8$.

Case (1.3.2.1.2): λ_1 moves at most 46 steps as γ is moving towards m_4 . Suppose that λ_1 moves ℓ steps towards v_4 , where $\ell \leq 46$. γ first moves to $L_{U,\ell+3}$; she then moves along the side path of $L_{U,\ell+3}$ parallel to B_9 until she reaches the corner of $L_{U,\ell+3}$ that is $92 - 2\ell$ edges away from v_4 . γ then moves to v_4 in $92 - 2\ell$ rounds. Since $d_{\mathcal{D}}(u_3, v_3) \geq 12$, γ can safely reach at least one of $\{o_4, o_5, q_4\}$ after reaching v_4 . Note that if γ moves to q_4 using the preceding strategy, then she requires a total of $202 + \ell$ rounds (starting at the round when she moves away from o). On the other hand, the cops need at least 196 rounds to reach U_9 , λ_3 needs at least 12 rounds to reach U_4 , and λ_1 needs at least 96 rounds to reach a neighbour of $U_4 \cup U_5 \cup U_9$. Thus if γ safely reaches q_4 in another 100 rounds, then Lemma 5.7 may be applied to $U_4 \cup U_5 \cup U_9$.

Case (1.3.2.2): $d_{\mathcal{D}}(u_3, v_3) \leq 11$.

Case (1.3.2.2.1): $d_{\mathcal{D}}(u_2, q_1) \geq 110$ and $d_{\mathcal{D}}(u_2, U_6) \geq 12$. γ first moves to v_1 in 98 rounds. If λ_3 is not in U_2 at the end of the 98-th round, then γ can safely reach o_2 in another 98 rounds. Suppose λ_3 does reach U_2 by the end the 98-th round, using up at least 89 turns in the process. After reaching v_1 , γ continues moving along B_1 until she reaches q_1 in another 100 rounds. Since $d_{\mathcal{D}}(u_2, q_1) \geq 110$ and $d_{\mathcal{D}}(u_3, q_1) \geq 199$, γ can safely reach q_1 . Furthermore, since $d_{\mathcal{D}}(u_2, U_6) \geq 12$, λ_1 needs at least 98 rounds to reach a neighbour of $U_1 \cup U_2 \cup U_6$ and λ_3 uses up at least 89 turns to reach U_2 , either γ may safely reach $u_1 \in U_2 \cup U_3$ and $u_2 \in U_3 \cup U_4$.

Case (1.3.2.2.2): $d_{\mathcal{D}}(u_2, q_1) \leq 109$ or $d_{\mathcal{D}}(u_2, U_6) \leq 11$. Then $d_{\mathcal{D}}(u_2, v_5) \geq 12$. γ may thus apply a strategy similar to that in Case (1.2.2.1), first moving towards m_5 and then either safely reaching o_5 or moving to one of $\{z_2, q_4\}$ and subsequently applying the strategy in Lemma 5.7 to either $U_5 \cup U_9 \cup U_{10}$ or $U_4 \cup U_5 \cup U_9$.

Case (2): There does not exist a corner $v \in \{v_1, v_2, v_3, v_4, v_5\}$ of $L_{U,49}$ such that $d_{\mathcal{D}}(u_1, v) \geq 98$, $d_{\mathcal{D}}(u_2, v) \geq 99$ and $d_{\mathcal{D}}(u_2, v) \geq 99$. Without loss of generality, assume that u_2 is in U_1 while u_3 is




Figure 15: An escape path of γ in Case (B.2.1.1.1.1).

in U_3 . Since $d_{\mathcal{D}}(u_1, v_4) \leq 98$, we have $d_{\mathcal{D}}(u_1, v_4) = 97$.

Case (2.1): $d_{\mathcal{D}}(u_3, v_3) \leq 11$. The following two cases are distinguished.

Case (2.1.1): $d_{\mathcal{D}}(u_2, v_5) \geq 12$. γ begins moving towards m_5 . We further distinguish two cases.

Case (2.1.1.1): λ_1 moves at least 47 steps as γ is moving towards m_5 . Suppose that as γ is approaching m_5 , λ_1 moves z steps for some $z \geq 47$. γ then continues moving until she reaches m_5 in 98 rounds. Note that λ_2 and λ_3 can move a total of at most 51 steps between the turn γ moves away from o and the turn after γ reaches m_5 . So λ_2 is at most 39 vertices closer to o_5 than γ is after γ reaches m_5 . We may assume that at least one of λ_1, λ_2 reaches U_5 just after γ reaches m_5 (otherwise, γ can safely reach o_5 in another 98 rounds).

Case (2.1.1.1.1): λ_1 reaches U_5 before λ_2 . Note that λ_2 is still at least 11 edges away from U_5 just after γ reaches m_5 . γ starts by moving towards o_5 until she reaches $L_{U_5,4}$; γ then moves along the path highlighted in Figure 15. An argument very similar to those used in earlier cases shows that either γ can move to o_{10} without being caught after reaching t_{19} , or γ can continue moving until she safely reaches z_2 , at which point Lemma 5.7 may be applied to $U_5 \cup U_9 \cup U_{10}$.

Case (2.1.1.1.2): λ_2 reaches U_5 before λ_1 . Suppose that λ_2 is ℓ vertices closer to o_5 than γ is during the turn after γ reaches m_5 . Note that $\ell \leq 39$. γ starts by moving towards o_5 . Suppose that as γ is approaching o_5 , λ_2 skips j turns. If $j > \ell$ then γ can safely reach o_5 . So assume that $j \leq \ell$. γ continues moving towards o_5 until she reaches $L_{U_5,4+\ell-j}$. She then moves along the path highlighted in Figure 16. One can verify that after reaching q_4 , either γ can safely reach o_9 in another 98 rounds, or Lemma 5.7 may be applied to $U_4 \cup U_5 \cup U_9$.

Case (2.1.1.2): λ_1 moves at most 46 steps as γ is moving towards m_5 . Suppose that λ_1 moves ℓ steps towards v_4 , where $\ell \leq 46$. γ first moves to $L_{U,\ell+3}$; she then moves along the side path of $L_{U,\ell+3}$ parallel B_{10} until she reaches the corner of $L_{U,\ell+3}$ that is $92 - 2\ell$ edges away from v_4 . γ then moves to v_4 in $92 - 2\ell$ rounds; note that λ_3 cannot catch γ just after γ reaches v_4 because he is at least 4 edges away from U_4 . Since $d_{\mathcal{D}}(u_2, v_5) \geq 12$, γ can safely reach either o_5 or q_4 after reaching v_4 . Note that if γ moves to q_4 using the preceding strategy, then she requires a total of $202 + \ell$




Figure 16: An escape path of γ in Case (B.2.1.1.1.2).




Figure 17: An escape path of γ in Case (B.2.1.2).

rounds (starting at the round when she moves away from o). On the other hand, the cops need at least 196 rounds to reach U_9 , λ_2 needs at least 12 rounds to reach U_5 , and λ_1 needs at least 96 rounds to reach a neighbour of $U_4 \cup U_5 \cup U_9$. Thus if γ can safely reach q_4 in another 100 rounds, then Lemma 5.7 may be applied to $U_4 \cup U_5 \cup U_9$.

Case (2.1.2): $d_{\mathcal{D}}(u_2, v_5) \leq 11$. Both $d_{\mathcal{D}}(u_2, v_5) \leq 11$ and $d_{\mathcal{D}}(u_3, v_3) \leq 11$ hold. γ starts moving towards m_2 . Note that at most one of λ_2 and λ_3 can reach U_2 before or just after γ reaches m_2 . We may assume that either λ_2 or λ_3 reaches U_2 before or just after γ reaches m_2 .

Suppose that λ_3 reaches U_2 before λ_2 . Suppose λ_3 is ℓ vertices closer to o_2 than γ is just after γ reaches m_2 . Note that $\ell \leq 9$. γ starts moving towards o_2 . Suppose λ_3 skips j turns as γ is approaching o_2 . If $j > \ell$, then γ can safely reach o_2 . Assume now that $j \leq \ell$. γ moves towards o_2 until she reaches $L_{U_2,4+\ell-j}$, continuing along the path highlighted in Figure 17 until she reaches q_1 . One may directly verify (in a way that is similar to earlier cases) that either γ can safely reach o_6 , or Lemma 5.7 may be applied to $U_1 \cup U_2 \cup U_6$. The case that λ_2 reaches U_2 before λ_3 may be handled similarly; in this case γ should move from t_{22} to t_2 instead.

Case (2.2): $d_{\mathcal{D}}(u_3, v_3) \geq 12$. Observe that this case is almost symmetrical to Case (2.1.1) and a parallel argument may be applied. More precisely, note that if one maps the set of corner vertices of U to itself as follows: $v_4 \to v_4, v_5 \to v_3, v_3 \to v_5, v_1 \to v_2, v_2 \to v_1$, and extend this mapping so as to obtain an automorphism σ of \mathcal{D} , then γ may apply a strategy similar to that in Case (2.1.1) for $\sigma(\mathcal{D})$ (with the appropriate transformed vertices).

C Detailed analysis of Algorithm 6 for Case (C)

Without loss of generality, assume that λ_2 is currently not in U while both λ_1 and λ_3 are currently in U. As the proof techniques in the present case are so similar to those in Cases (A) and (B), we will omit many proof details and refer to strategies for γ in previous cases.

Case (1): There is at least one corner v_i of $L_{U,49}$ such that $d_{\mathcal{D}}(v_i, u') \geq 99$ for all $u' \in \{u_2, u_3\}$ and $d_{\mathcal{D}}(v_i, u_1) \geq 98$. We first assume that i = 1. As in Case (B), define $F := U_{10} \cup U_6 \cup U_1 \cup U_2 \cup U_7$.

Case (1.1): u_2 is in F. First, suppose that $d_{\mathcal{D}}(u_2, v_5) \leq 98$. (This implies that u_2 is in U_1 .) If $d_{\mathcal{D}}(u_3, B_{10}) \leq 98$, then γ moves to v_2 in 98 rounds; Lemma 5.6 may then be applied to $U_2 \cup U_3$. Now suppose that $d_{\mathcal{D}}(u_3, B_{10}) \geq 99$. If $d_{\mathcal{D}}(u_2, v_5) \geq 12$, then γ may apply a winning strategy similar to that in Case (B.2.1.1). Now suppose that $d_{\mathcal{D}}(u_2, v_5) \leq 11$. If $d_{\mathcal{D}}(u_3, B_7) \leq 50$, then γ may apply the winning strategy in Lemma 5.5, first moving to p in 98 rounds. Now suppose $d_{\mathcal{D}}(u_3, B_7) \geq 51$. γ first moves to v_1 in 98 rounds. Note that $d_{\mathcal{D}}(u_2, q_1) \geq 185$. If λ_3 is not in U_2 just after the round when γ reaches v_1 , then γ can safely reach o_2 in another 98 rounds. If λ_3 is in U_2 just after the 98-th round. Thus λ_2 is at least 138 edges away from q_1 just after the 98-th round. γ can now safely move to q_1 in 100 rounds, and then to o_6 in another 98 vertices.

Second, suppose that $d_{\mathcal{D}}(u_2, v_5) \geq 99$. We distinguish the following cases.

Case (1.1.1): $d_{\mathcal{D}}(u_3, v_5) \leq 101$, $d_{\mathcal{D}}(u_3, v_4) \leq 101$ and for all $i \in \{1, 2, 3\}$, $d_{\mathcal{D}}(u_3, v_i) \geq 100$. First, suppose that $d_{\mathcal{D}}(u_2, v_2) \leq 98$. If $d_{\mathcal{D}}(u_3, U_4) + d_{\mathcal{D}}(u_2, U_3) \geq 5$, then γ first moves to v_3 in 98 rounds. If neither λ_1 nor λ_3 reaches U_4 at the end of the 98-th round, then γ can move to o_4 without being caught in another 98 rounds. If either λ_1 or λ_3 reaches U_4 at the end of the 98-th round (or if both λ_1 and λ_3 reach U_4 at the end of the 98-th round), then γ can safely reach q_3 in another 100 rounds. After reaching q_3 , either γ can safely reach o_8 in another 98 rounds, or Lemma 5.7 may be applied to $U_3 \cup U_4 \cup U_8$.

If $d_{\mathcal{D}}(u_3, U_4) + d_{\mathcal{D}}(u_2, U_3) \leq 4$, then γ first moves to v_1 in 98 rounds. If λ_3 does not reach U_1 at the end of the 98-th round, then γ can safely reach o_1 in another 98 rounds. If λ_3 reaches U_1 at the end of the 98-th round, then, since $d_{\mathcal{D}}(u_2, q_1) + d_{\mathcal{D}}(u_3, U_1) \geq 100 - d_{\mathcal{D}}(u_3, U_4) + 196 - d_{\mathcal{D}}(u_2, U_3) \geq 288$, γ can move safely towards q_1 in another 100 rounds, and then safely reach o_6 using an additional 98 rounds.

Second, suppose $d_{\mathcal{D}}(u_2, v_2) \geq 99$. Suppose $u_2 \in V(U_7)$. If $u_2 \neq q_2$, then γ first moves to v_2 in 98 rounds. If λ_2 does not reach B_2 by the end of the 98-th round, then γ can safely reach either o_2 or o_3 . If λ_2 does reach B_2 by the end of the 98-th round, then both λ_1 and λ_3 are still at least 2 edges away from $U_2 \cup U_3$ at the end of the 98-th round, and therefore Lemma 5.6 may be applied to

 $U_2 \cup U_3$. Suppose $u_2 = q_2$. If $d_{\mathcal{D}}(u_3, U_4) \geq 4$, then γ first moves to v_3 in 98 rounds. If neither λ_1 nor λ_3 is in U_4 at the end of the 98-th round, then γ can safely reach o_4 in another rounds. If either λ_1 or λ_3 is in U_4 at the end of the 98-th round, then γ continues moving along B_3 until she reaches q_3 using another 100 rounds. Then either γ can safely move to o_8 in another 98 rounds, or Lemma 5.7 may be applied to $U_3 \cup U_4 \cup U_8$. If $d_{\mathcal{D}}(u_3, U_4) \leq 3$, then γ first moves to v_1 in 98 rounds. After reaching v_1 , γ can either safely reach o_1 using another 98 rounds, or γ can move to q_1 in another 100 rounds and then apply the strategy in Lemma 5.7 to $U_1 \cup U_2 \cup U_6$.

Now suppose $u_2 \notin V(U_7)$. If $d_{\mathcal{D}}(u_3, U_4) + d_{\mathcal{D}}(u_2, U_3) \geq 3$, then γ first moves to v_3 in 98 rounds. If U_3 (resp. U_4) does not contain any cop at the end of the 98-th round, then γ safely moves to o_3 (resp. o_4) using another 98 rounds. Suppose each of U_3 and U_4 contains a cop at the end of the 98-th round. Since $d_{\mathcal{D}}(u_i, U_4) + d_{\mathcal{D}}(u_2, U_3) \geq 3$ whenever $i \in \{1, 3\}$, it follows that at the end of the 98-th round, γ can safely move along B_3 to q_3 using another 100 rounds. After reaching q_3 , γ can either safely reach o_8 using another 98 rounds or apply the strategy in Lemma 5.7 to $U_3 \cup U_4 \cup U_8$. If $d_{\mathcal{D}}(u_3, U_4) + d_{\mathcal{D}}(u_2, U_3) \leq 2$, then γ first moves to v_1 in 98 rounds. If no cop is in U_1 at the end of the 98-th round, then γ can safely reach o_1 in another 98 rounds; otherwise, γ can move along B_1 and reach q_1 without being caught; she can then move safely to o_6 in another 98 rounds.

Case (1.1.2): $d_{\mathcal{D}}(u_3, v_3) \leq 101$, $d_{\mathcal{D}}(u_3, v_4) \leq 101$ and for all $i \in \{1, 2, 5\}$, $d_{\mathcal{D}}(u_3, v_i) \geq 100$. Note that u_3 and u_1 are each at least 99 edges away from $U_1 \cup U_2$. First, suppose that $u_2 \neq q_1$. γ moves to v_1 in 98 rounds. If λ_2 does not move between the 1-st and the 98-th round, then some $U_i \in \{U_1, U_2\}$ does not contain any cop at the end of the 98-th round. γ may then safely reach o_i in another 98 rounds. If λ_2 moves at least one step between the 1-st and the 98-th round, then both λ_3 and λ_1 are each at least 2 edges away from $U_1 \cup U_2$ at the end of the 98-th round. One may then apply Lemma 5.6 to $U_1 \cup U_2$.

Second, suppose that $u_2 = q_1$. If $d_{\mathcal{D}}(u_3, B_{10}) \geq 50$, then γ moves to v_5 in 98 rounds. At the end of the 98-th round, γ can either safely reach o_5 in another 98 rounds, or move towards q_5 and then to o_{10} in another 198 rounds. If $d_{\mathcal{D}}(u_3, B_8) \geq 50$, then γ moves to v_2 in 98 rounds. An argument similar to that in the preceding case (that is, when $d_{\mathcal{D}}(u_3, B_{10}) \geq 50$) shows that γ can either safely reach o_3 in another 98 rounds or safely reach o_7 in another 198 rounds.

Case (1.1.3): $d_{\mathcal{D}}(u_3, v_2) \leq 101$, $d_{\mathcal{D}}(u_3, v_3) \leq 101$ and for all $i \in \{1, 4, 5\}$, $d_{\mathcal{D}}(u_3, v_i) \geq 100$. γ moves to v_5 in 98 rounds. An argument very similar to that in Case (1.1.2) shows that γ can either safely reach o_5 in another 98 rounds, or safely reach o_{10} in another 198 rounds.

Case (1.1.4): $d_{\mathcal{D}}(u_3, v_1) \leq 101$, $d_{\mathcal{D}}(u_3, v_2) \leq 101$ and for all $i \in \{3, 4, 5\}$, $d_{\mathcal{D}}(u_3, v_i) \geq 100$. γ pursues the same winning strategy as that in Case (1.1.3).

Case (1.1.5): $d_{\mathcal{D}}(u_3, v_1) \leq 101$, $d_{\mathcal{D}}(u_3, v_5) \leq 101$ and for all $i \in \{2, 3, 4\}$, $d_{\mathcal{D}}(u_3, v_i) \geq 100$. First, suppose that $d_{\mathcal{D}}(u_2, v_2) \leq 98$. γ moves to v_3 in 98 rounds. If neither λ_1 nor λ_3 is in U_4 at the end of the 98-th round, then γ may safely reach o_4 in another 98 rounds. If either λ_1 or λ_3 is in U_4 at the end of the 98-th round, then λ_2 must still be at least 195 edges away from q_3 at the end of the 98-th round. Thus γ may safely move to q_3 in 100 rounds, and then to o_8 in another 98 rounds.

Second, suppose that $d_{\mathcal{D}}(u_2, v_2) \geq 99$. If $u_2 = q_2$, then γ employs the winning strategy in the preceding case (that is, the case when $d_{\mathcal{D}}(u_2, v_2) \leq 98$). If $u_2 \neq q_2$, then γ moves to v_2 in 98 rounds. If λ_2 moves at least one step between the 1-st and the 98-th round, then λ_1 and λ_3 are each at least 2 edges away from $U_2 \cup U_3$ at the end of the 98-th round (note that since $d_{\mathcal{D}}(u_3, v_1) \geq 99$ by the

case assumption and v_1 is the vertex of U_2 that is closest to u_3 , $d_{\mathcal{D}}(u_3, U_2) \geq 99$) and therefore Lemma 5.6 may be applied to $U_2 \cup U_3$. If λ_2 does not move between the 1-st and the 98-th round, then there is some $U_i \in \{U_2, U_3\}$ such that U_i does not contain any cop at the end of the 98-th round. Thus γ may safely reach o_i in another 98 rounds.

Case (1.2): u_2 is not in F. First, suppose that $d_{\mathcal{D}}(u_2, U_1 \cup U_2) \geq 3$ or $d_{\mathcal{D}}(u_3, U_1 \cup U_2) \geq 3$. γ first moves to v_1 in 98 rounds. If some $U_i \in \{U_1, U_2\}$ does not contain any cop at the end of the 98-th round, then γ moves to o_i in another 98 rounds. If both U_1 and U_2 contain at least one cop at the end of the 98-th round, then γ continues moving towards q_1 . Since each cop requires at least 196 rounds (from his starting position) to reach q_1 but at least 2 cops need more than 2 rounds to reach $U_1 \cup U_2$ (and no cop can reach v_1 in 98 rounds), γ can safely get from v_1 to q_1 in 100 rounds, and then move from q_1 to o_6 in another 98 rounds.

Second, suppose that $d_{\mathcal{D}}(u_2, U_1 \cup U_2) \leq 2$ and $d_{\mathcal{D}}(u_3, U_1 \cup U_2) \leq 2$. γ then moves to p in 98 rounds. One may then apply Lemma 5.5 to obtain a winning strategy for γ .

- Case (2): For each corner v' of $L_{U,49}$, it holds that $d_{\mathcal{D}}(v',u') \leq 98$ for some $u' \in \{u_2,u_3\}$ or $d_{\mathcal{D}}(v',u_1) \leq 97$ (or both inequalities hold).
- Case (2.1): $d_{\mathcal{D}}(u_2, v_2) \leq 98$, $d_{\mathcal{D}}(u_2, v_3) \leq 98$, $d_{\mathcal{D}}(u_3, v_1) \leq 98$, $d_{\mathcal{D}}(u_3, v_5) \leq 98$ and $d_{\mathcal{D}}(u_1, v_4) = 97$. Suppose that $d_{\mathcal{D}}(u_2, v_3) \geq 12$. γ may then apply the winning strategy in Case (B.2.2). Now suppose that $d_{\mathcal{D}}(u_2, v_3) \leq 11$. Then $d_{\mathcal{D}}(u_2, U_2) \geq 89$ and $d_{\mathcal{D}}(u_2, m_2) \geq 139$. Consider the following case distinction: (i) $d_{\mathcal{D}}(u_3, m_2) \leq 98$ and (ii) $d_{\mathcal{D}}(u_3, m_2) \geq 99$.
- (i) Notice that in this case, $d_{\mathcal{D}}(u_3, m_5) \geq 99$ and $d_{\mathcal{D}}(u_3, v_5) \geq 49$. γ may then apply a winning strategy similar to that in Case (B.2.1.1).
- (ii) It follows from the inequalities $d_{\mathcal{D}}(u_3, m_2) \geq 99$ and $d_{\mathcal{D}}(u_3, v_5) \leq 98, d_{\mathcal{D}}(u_3, v_1) \leq 98$ that $d_{\mathcal{D}}(u_3, v_1) \geq 49$. γ may then apply the winning strategy in Lemma 5.5, first moving to m_2 in 98 rounds (note that the winning strategy applies in this case even though λ_2 is not in U at the start).
- Case (2.2): $d_{\mathcal{D}}(u_2, v_1), d_{\mathcal{D}}(u_2, v_5) \leq 98, d_{\mathcal{D}}(u_3, v_2), d_{\mathcal{D}}(u_3, v_3) \leq 98$ and $d_{\mathcal{D}}(u_1, v_4) = 97$. As in Case (B.2.1), we first suppose that $d_{\mathcal{D}}(u_2, v_5) \geq 12$. γ may then employ the winning strategy in Case (B.2.1.1). Now suppose that $d_{\mathcal{D}}(u_2, v_5) \leq 11$. Then $d_{\mathcal{D}}(u_2, m_2) \geq 139$ and $d_{\mathcal{D}}(u_2, v_1) \geq 89$.
- (i) $d_{\mathcal{D}}(u_3, m_2) \geq 99$. Then $d_{\mathcal{D}}(u_3, v_2) \geq 49$. γ moves to m_2 in 98 rounds, employing the winning strategy in Lemma 5.5.
- (ii) $d_{\mathcal{D}}(u_3, m_2) \leq 98$. Then $d_{\mathcal{D}}(u_3, m_4) \geq 99$. γ moves to m_4 in 98 rounds, employing the winning strategy in Case (B.2.2).